



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, 15/19, C07K 14/715, 14/52, C12Q 1/68, G01N 33/53, 33/68, C07K 16/24, 16/28	A2	(11) International Publication Number: WO 99/33980 (43) International Publication Date: 8 July 1999 (08.07.99)
(21) International Application Number: PCT/US98/27474 (22) International Filing Date: 22 December 1998 (22.12.98) (30) Priority Data: 60/068,959 30 December 1997 (30.12.97) US 09/212,270 16 December 1998 (16.12.98) US (71) Applicant: CHIRON CORPORATION [US/US]; 4560 Horton Street -R440, Emeryville, CA 94608 (US). (72) Inventors: TRIBOULEY, Catherine; Chiron Corporation, In- tellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US). POT, David; Chiron Corporation, Intel- lectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US). KASSAM, Altaf; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US). LAMSON, George; Chiron Cor- poration, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US). (74) Agents: POTTER, Jane, E., R. et al.; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished</i> <i>upon receipt of that report.</i>
(54) Title: MEMBERS OF TNF AND TNFR FAMILIES (57) Abstract <p>Members of the TNF and the TNFR superfamily of proteins have been identified. These proteins are promising targets for therapeutic intervention and mimesis. TNF-L and TNFR-L proteins can be used to induce cell death and/or proliferation of cells. Members of these superfamilies have been implicated in a broad variety of disease processes, making them central biological and physiological regulators.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

MEMBERS OF TNF AND TNFR FAMILIES

5

BACKGROUND OF THE INVENTION

Tumor necrosis factor (TNF) is a pro-inflammatory cytokine which is produced by a wide spectrum of cells. It has a key role in host defense and immunosurveillance, mediating complex cellular responses. In excess, TNF may have detrimental effects.

10 Two specific, high affinity cell surface receptors, p55 TNF-R and p75 TNF-R, function as transducing elements, providing the intracellular signal for cell responses to TNF. While both types of TNF receptors are expressed by almost all cell types, the p75 receptor has been shown to be expressed primarily by cells of the immune system (B and T cells), cells of myeloid origin, and endothelial cells. Both receptors participate in
15 the induction of NF κ B and interleukin-6, in the generation of lymphocyte activated killer (LAK) cells, and in the proliferation of natural killer (NK) cells, as well as in anti-proliferation, cytotoxicity, and apoptosis.

TNF signaling to cells is largely mediated by the p55 TNF-R, while the main function of the p75 surface receptor is "ligand passing," *i.e.*, TNF presentation to the p55
20 TNF-R. Presence of the cell surface p75 TNF receptor greatly enhances the rate of association of TNF to the p55 TNF receptor and may reverse the desensitization of p55 TNF-R to TNF. Pharmaceutical agents which affect p75 TNF-R may have a general impact on TNF function, including those activities in which the major signaling receptor is the p55 TNF-R.

25 The TNF-Rs also mediate many non-overlapping functions: the p55 receptor is involved in interleukin-2 receptor induction, anti-viral activities, growth stimulation, HLA antigen expression, and endothelial cell adhesion, while the p75 receptor mediates the TNF-induced thymocyte proliferation.

The p55 and p75 TNF-Rs are members of a superfamily which includes nerve
30 growth factor receptor (NGFR), Fas antigen, CD27, CD30, CD40, OX40 and 4-1BB. The cysteine-rich domains of the extracellular part of these receptors are homologous to

several viral proteins produced by cowpox virus, Shope fibroma virus, and the myxoma virus.

Because of the central role of TNF and its receptors in host defense and immunosurveillance, there is a need in the art to identify new members of the TNF and
5 TNFR superfamilies.

SUMMARY OF THE INVENTION

It is an object of the invention to provide new members of the TNF and TNFR families, as well as methods of screening for compounds capable of modifying the
10 activities of these proteins. This and other objects of the invention are provided by one or more of the embodiments described below.

One embodiment of the invention is an isolated human protein having an amino acid sequence which is at least 85% identical to an amino acid sequence selected from the group consisting of SEQ ID NOS:1 and 2. Percent identity is determined using a
15 Smith-Waterman homology search algorithm using an affine gap search with a gap open penalty of 12 and a gap extension penalty of 1.

Another embodiment of the invention is a fusion protein comprising a first protein segment and a second protein segment fused together by means of a peptide bond. The first protein segment consists of a protein having an amino acid sequence
20 selected from the group consisting of SEQ ID NOS:1 and 2.

Still another embodiment of the invention is a preparation of antibodies which specifically bind to a protein having an amino acid sequence selected from the group consisting of SEQ ID NOS:1 and 2.

Even another embodiment of the invention is a cDNA molecule which encodes a
25 protein having an amino acid sequence which is at least 85% identical to an amino acid sequence selected from the group consisting of SEQ ID NOS:1 and 2. Percent identity is determined using a Smith-Waterman homology search algorithm using an affine gap search with a gap open penalty of 12 and a gap extension penalty of 1.

Yet another embodiment of the invention is a cDNA molecule which is at least
30 85% identical to a nucleotide sequence selected from the group consisting of SEQ ID

NOS:6 and 7. Percent identity is determined using a Smith-Waterman homology search algorithm using an affine gap search with a gap open penalty of 12 and a gap extension penalty of 1.

5 A further embodiment of the invention is an isolated and purified subgenomic polynucleotide comprising a nucleotide sequence which hybridizes to a nucleotide sequence selected from the group consisting of SEQ ID NOS:6 and 7 after washing with 0.2X SSC at 65 °C. The nucleotide sequence encodes a protein having an amino acid sequence selected from the group consisting of SEQ ID NOS:1 and 2.

10 Another embodiment of the invention is a construct comprising a promoter and a polynucleotide segment encoding an amino acid sequence selected from the group consisting of SEQ ID NOS:1 and 2. The polynucleotide segment is located downstream from the promoter. Transcription of the polynucleotide segment initiates at the promoter.

15 Still another embodiment of the invention is a host cell comprising a construct which comprises a promoter and a polynucleotide segment encoding an amino acid sequence selected from the group consisting of SEQ ID NOS:1 and 2.

Even another embodiment of the invention is a recombinant host cell comprising a new transcription initiation unit, wherein the new transcription initiation unit comprises in 5' to 3' order (a) an exogenous regulatory sequence, (b) an exogenous exon, and (c) a splice donor site. The new transcription initiation unit is located upstream of a coding sequence of a gene. The coding sequence is selected from the group consisting of SEQ ID NOS:6 and 7. The exogenous regulatory sequence controls transcription of the coding sequence of the gene.

20 Yet another embodiment of the invention is a method of screening for a compound capable of modulating cell death inducing activity of a protein. A first population of cells and a protein are incubated in the presence of a test compound. The protein comprises an amino acid sequence selected from the group of amino acid sequences shown in SEQ ID NOS:1-5. A second population of cells and the protein are incubated in the absence of a test compound. Viability of the first and second populations is determined. A test compound which increases or decreases viability of the first population relative to the second population is identified as capable of

30

modulating the cell death inducing activity of the protein.

Even another embodiment of the invention is a method of identifying a binding partner of a first protein. A first protein is incubated with a second protein. The first protein comprises an amino acid sequence selected from the group consisting of SEQ ID
5 NOS:1-5. Formation of a complex between the first and second proteins is detected. Formation of the complex identifies the second protein as a binding partner of the first protein.

The present invention thus provides the art with the amino acid sequences of proteins which are new members of the TNF and TNFR families, as well as nucleotide
10 sequences of polynucleotides which encode the proteins. These proteins and polynucleotides can be used to enhance or decrease TNF activities thereby providing therapeutic benefits, such as induction of cell death, lymphoid organogenesis, or host bacterial resistance, and inhibition of endotoxic shock, contact hypersensitivity, delayed type hypersensitivity, or immunocompetence of a transplant recipient. Methods of
15 diagnosing neoplasia or predisposition to neoplasia are also provided. Proteins of the present invention are also useful for identifying compounds which can regulate the TNF-like or TNFR-like activities of these proteins.

BRIEF DESCRIPTION OF THE DRAWINGS

20 **Figure 1.** Figure 1A shows the protein sequence of human TNFL1. The transmembrane domain is underlined. Figure 1B shows alignments of the extracellular domains of several members of the TNF family with TNFL1. The conserved amino acids are boxed.

Figure 2. Figure 2A shows *TNFL1* mRNA expression in human tissues and cell
25 lines. Figure 2B shows *TNFL1* expression in mouse tissues. PBL, peripheral blood lymphocytes; HL60 promyelocytic leukemia HL60; HeLa, HeLa cell line S3; K562, chronic myelogenous leukemia K562; MOLT-4, lymphoblastic leukemia MOLT-4; Raji, Burkitt's lymphoma Raji; SW480, colorectal adenocarcinoma SW480; A549, lung carcinoma A549; G361, melanoma G361.

30 **Figure 3.** Figure 3 shows detection of TNFL1 expression using a purified

polyclonal anti-peptide antibody.

Figure 3A shows a Western blot analysis using the affinity-purified D2710 antibody at a final concentration of 1 µg/ml. The extracts analyzed were total cell extracts of human monocytes isolated from PBMCs (lanes 1 and 2) and mouse bone marrow-derived dendritic cells (lanes 3 and 4). Lanes 5 and 6 contained 20 ng of TNFL1 protein purified from *E. coli*. In lanes 2, 4, and 6 the antibody was pre-incubated with a hundred-fold molar excess of the peptide used to generate the antibody.

Figure 3B shows intracellular staining of baculovirus-infected cells expressing TNFL1, using the D2710 antibody or a control rabbit antibody at a concentration of 10 µg/ml. The insect cells were fixed and permeabilized before staining and flow cytometry analysis.

Figure 3C shows the pattern of expression of TNFL1 in mouse spleen sections. Left panels: rabbit antibody control plus secondary antibody; right panels, anti-TNFL1 antibody plus secondary antibody. The magnification is 100x in the top panels and 200x in the bottom panels. RP, red pulp; PALS, periarteriolar lymph sheath.

Figure 3D shows staining of adjacent murine spleen sections for Thy-1.2, B220, TNFL1, CD11c, and Mac-3.

Figure 4. Figure 4 shows flow cytometric analysis of the cell-surface expression of TNFL1 on sub-populations of human PBMCs and cultured cells.

Figure 4A shows expression of TNFL1 on CD14+, CD19+, CD4+, and CD8+ human PBMCs and on mouse bone marrow-derived dendritic cells. Human PBMCs were stained with affinity purified D2710, followed by FITC-conjugated anti-rabbit IgG. They were subsequently stained with PE-conjugated CD14, CD19, CD4, or CD8. The histograms show TNFL1 expression (FITC) on PE-positive gated cells. The dotted lines represent the staining in the absence of primary antibody D2710.

Figure 4B shows upregulation of TNFL1 surface expression on T cells after activation with anti-CD3 and anti-CD28 at 10 µg/ml in the presence of IL2 at 50 µg/ml for six days.

Figure 5. Figure 5 shows the effect of recombinant TNFL1 on activated T and B cells.

Figure 5A shows the amino acid sequence of a soluble form of TNFL1 (sequence enclosed in brackets) which was expressed in *E. coli* as a fusion protein and used for all biological assays. The cleavage sites identified after microsequencing of the truncated form of the protein are represented as vertical bars.

5 Figure 5B shows inhibition of DNA synthesis in activated but not in resting T cells by TNFL1. T cells enriched from human PBMCs were either activated with 10 µg/ml of anti-CD3 and 10 µg/ml of anti-CD28 (left panel) or untreated (right panel) for 24 hours. TNFL1 was then added at various concentrations, and ³H-thymidine incorporation over a period of 8 hours was measured 48 hours after addition of TNFL1.

10 Figure 5C shows inhibition of DNA synthesis by TNFL1 in activated B cells and activated T cells. B cells were activated with 10 µg/ml of anti-CD40 antibody. T cells were activated as described in Figure 5B. TNFL1 was added 48 hours after activation. ³H-thymidine incorporation over a period of 8 hours was measured 48 hours after addition of TNFL1.

15 Figure 5D shows that TNFL1 induces apoptosis of activated T cells. T cells were activated with anti-CD3 and anti-CD28 for 48 hours. TNFL1 was added at 2 µg/ml and incubated for an additional 48 hours. Apoptosis was assessed by TUNEL assay. (R1), blasting cells; (R2), apoptotic cells; (R3), resting non-apoptotic cells.

20 Figure 5E shows that TNFL1 induces NFκB activation in Jurkat cells. An electrophoretic mobility shift assay was performed with an NFκB probe on nuclear cell extracts prepared from Jurkat cells treated with PMA (1 µg/ml) or TNFL1 (3 µg/ml) for one hour. Wild-type (wt, 20 ng) and mutated (mut, 100 ng) non-radiolabeled oligonucleotides were used as competitors in the reaction.

25

DETAILED DESCRIPTION OF THE INVENTION

New members of the TNF and TNFR families are a discovery of the present invention. In particular, the present invention identifies two cDNA clones which encode new members of TNF ligand family and three cDNA clones which encode new members of TNFR family. Proteins and polynucleotides of the invention provide the art with diagnostic and therapeutic reagents as well as tools for discovering other therapeutic agents.

Two cDNA clones have been identified which show significant homology to members of TNF family. One cDNA clone (SEQ ID NO:6) was isolated from a human liver cDNA library. It encodes a protein designated as TNFL1 (SEQ ID NO:1), which has 27%-30% homology with TNF and 38% homology with lymphotoxin. TNFL1 contains a transmembrane domain. The *TNFL1* cDNA clone encodes a 3 kb mRNA which can be detected in tissues associated with the immune system, such as peripheral blood lymphocytes, spleen, and thymus, as well as small intestine and ovary. TNFL1 protein is constitutively expressed on monocytes and B cells isolated from human peripheral blood lymphocytes, as well as on mouse dendritic cells and in mouse spleen. Expression of the protein can be up-regulated in natural killer cells. Activation of dendritic cells, for example with anti-CD4d antibody, can down-regulate expression of TNFL1 protein.

TNFL1 shares some common features with other members of the TNF family. TNFL1 is upregulated on activated T cells, as are TNF, Fas ligand, and CD30 ligand. TNFL1 induces activation of NF κ B, which is also triggered by every member of the TNF-R family with the exception of Fas and DR4. TNFL1 also leads to apoptosis of activated T cells, a well documented effect in the case of Fas ligand and TNF. TNFL1 differs in its expression pattern, however, when compared to TNF or Fas ligand. Fas ligand, which is involved in activation-induced cell death (AICD) of CD4+ T cells and tolerance to self-antigens, is classically expressed on activated T cells and the immuno-privileged eye and testis. TNF is constitutively expressed on both mature and immature thymocytes, is upregulated on activated T cells, and is induced by LPS on macrophages. In contrast, TNFL1 is constitutively expressed on antigen presenting cells,

specialized or not, such as monocytes, B cells from peripheral blood lymphocytes in humans, and likely on macrophages or dendritic cells in the red pulp and marginal zone of mouse spleen and cultured dendritic cells.

This localization in the blood and in the spleen suggests a possible function for
5 TNFL1 in the recognition process of blood-borne pathogens, such as bacteria or viruses. Furthermore, TNFL1 is expressed at the surface of dendritic cells cultured from bone marrow in the presence of GM-CSF. TNFL1 may therefore be expressed on a subtype of dendritic cells which were recently individualized, myeloid dendritic cells (MDCs), rather than on lymphoid dendritic cells (LDCs). MDCs share a precursor with
10 macrophages, are GM-CSF dependant, and are present in the marginal zone of secondary lymphoid tissues. LDCs, in contrast, are located in the T-cell zone of the secondary organs, are IL3-dependent, and share a precursor with T and B cells.

A second cDNA clone was isolated from an oligodT-primed library of a human ovarian tumor. It encodes a protein designated as TNFL2, which has about 25%
15 homology with TNF. TNFL2 does not have a transmembrane domain and thus can be secreted. The sequence of TNFL2 is shown in SEQ ID NO:5. The *TNFL2* cDNA clone detects a major population of mRNA in a range of about 1.5 kb to about 2 kb in tissues associated with the immune system, *e.g.*, peripheral blood lymphocytes and spleen. A slightly bigger mRNA is also expressed in spleen as well as in colon, prostate, and to a
20 lesser extent in ovary and small intestine. The full length polynucleotide sequence of *TNFL2* cDNA is shown in SEQ ID NO:10.

TNFL1 might be able to bind to the TNF receptors or to Fas, as a homotrimer or in association with another member of the TNF family. TNFL1 and TNFL2 may form heterodimers and work together in a manner similar to that of lymphotoxin α and β . The
25 TNF-like ligands disclosed herein can be used, *inter alia*, to induce cell death in tumors, to induce apoptosis of activated T cells, to induce inflammation, and to rescue resting T cells from apoptosis.

Proteins which are members of the TNFR superfamily have also been identified. These are soluble receptors which have the amino acid sequences shown in SEQ ID
30 NOS:2 and 3 (human) and SEQ ID NO:4 (mouse). These proteins are encoded by the

nucleotide sequences shown in SEQ ID NOS:7, 8, and 9, respectively. These receptors can be used, *inter alia*, to regulate the function of a TNF ligand which plays a rôle in apoptosis, inflammation, differentiation, or proliferation. Expression of the receptors can also be useful as markers for cancer, especially for colon cancer. Diseases which can be treated using the ligands and/or receptors of the TNF/TNFR superfamily include

The human TNF-like (TNF-L) and mammalian TNF receptor-like (TNFR-L) proteins or polypeptides, biologically active polypeptides or protein variants, and fusion proteins disclosed herein can be used in various therapeutic compositions and methods, as described below. Any naturally occurring variants of SEQ ID NOS:1-5 which may occur in human or mammalian tissues and which retain the functional properties of the TNF-L or TNFR-L proteins disclosed herein are biologically active TNF-L or TNFR-L variants. Non-naturally occurring TNF-L or TNFR-L variants which contain conservative amino acid substitutions relative to SEQ ID NOS:1-5 but which retain substantially the same ligand or receptor activity as naturally occurring TNF-L or TNFR-L are also biologically active TNF-L or TNFR-L variants.

Naturally or non-naturally occurring TNF-L or TNFR-L variants preferably are at least 85%, 90%, or 95% identical to SEQ ID NOS:1-5 and have similar biological functions, which are described below. More preferably, the molecules are 98% or 99% identical. Percent identity is determined using the Smith-Waterman homology search algorithm, using an affine gap search with a gap open penalty of 12 and a gap extension penalty of 1. The Smith-Waterman homology search algorithm is taught in Smith and Waterman, *Adv. Appl. Math.* (1981) 2:482-489.

Biologically active TNF-L or TNFR-L variants include glycosylated forms of the proteins, aggregative conjugates of the proteins with other molecules, and covalent conjugates of the proteins with unrelated chemical moieties. Covalent conjugates are prepared by linkage of functionalities to groups which are found in the amino acid chain or at the N- or C-terminal residues of the proteins of the invention by means well known in the art. TNF-L or TNFR-L variants also include allelic variants, species variants, and muteins. Truncations or deletions of regions which do not affect the biological functions

of the TNF-L or TNFR-L proteins disclosed herein are also biologically active TNF-L or TNFR-L variants.

A subset of mutants, called muteins, is a group of polypeptides with the non-disulfide bond participating cysteines substituted with a neutral amino acid, generally, with serines. These mutants may be stable over a broader temperature range than naturally occurring TNF-L or TNFR-L proteins. See Mark *et al.*, U.S. Pat. No. 4,959,314.

Biologically active TNF-L or TNFR-L polypeptides can comprise at least 6, 7, 8, 9, 10, 12, 15, 20, 25, 30, 35, 40, 50, 75, 100, 125, 150, 175, 200, 225, 250, or 275 contiguous amino acids of SEQ ID NO:1, at least 6, 7, 8, 9, 10, 12, 15, 20, 25, 30, 35, 40, 50, 75, 100, 125, or 150 contiguous amino acids of SEQ ID NO:2, at least 6, 7, 8, 9, 10, 12, 15, 20, 25, 30, 35, 40, 50, 75, 100, 125, 150, 175, or 200 contiguous amino acids of SEQ ID NO:3, at least 6, 7, 8, 9, 10, 12, 15, 20, 25, 30, 35, 40, 50, 75, 100, 125, 130, or 140 contiguous amino acids of SEQ ID NO:4, or at least 6, 7, 8, 9, 10, 12, 15, 20, 25, 30, 35, 40, 50, 75, 100, 125, 150, 175, or 200 contiguous amino acids of SEQ ID NO:5.

Polypeptide molecules having substantially the same amino acid sequences as the TNF-L or TNFR-L proteins disclosed herein but possessing minor amino acid substitutions which do not substantially affect the ability of the TNF-L or TNFR-L polypeptides to interact with their respective receptors or ligands are within the definition of biologically active TNF-L or TNFR-L polypeptide variants.

Preferably, biologically active TNF-L or TNFR-L polypeptides or polypeptide variants are at least 65%, 75%, 85%, 90%, 95%, 98%, or 99% identical to TNF-L or TNFR-L polypeptide fragments of SEQ ID NOS:1-5. Percent identity of potential polypeptides or polypeptide variants with fragments of SEQ ID NOS:1-5 is determined as described above.

Guidance in determining which amino acid residues can be substituted, inserted, or deleted without abolishing biological or immunological activity can be found using computer programs well known in the art, such as DNASTAR software. Preferably the amino acid changes in TNF-L or TNFR-L protein or polypeptide variants are conservative amino acid changes, *i.e.*, changes of similarly charged or uncharged amino

acids. Conservative replacements are those which take place within a family of amino acids which are related in their side chains. Genetically encoded amino acids are generally divided into four families: acidic (aspartate, glutamate); basic (lysine, arginine, histidine); non-polar (alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan); and uncharged polar (glycine, asparagine, glutamine, cystine, serine, threonine, tyrosine). Phenylalanine, tryptophan, and tyrosine are sometimes classified jointly as aromatic amino acids.

It is reasonable to expect that an isolated replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid will not have a major effect on the binding properties of the resulting TNF-L or TNFR-L molecule, especially if the replacement does not involve an amino acid at a binding site involved in an interaction of a TNF-L protein with its receptor or a TNFR-L protein with its ligand. Binding between a TNF-L protein and its receptor or a TNFR-L protein and its ligand can be measured, for example, using a yeast two-hybrid assay, as is known in the art (Fields & Song, *Nature* 340:245-46, 1989).

Alternatively, the amino acid sequence of a TNF-L or TNFR-L protein can be modified to alter its biological activity. For example, amino acids 174-193 (the TNF ligand binding domain) can be deleted in a TNFR-L protein to form an inactive variant of the TNFR-L protein and thereby inhibit or decrease the function of its ligand.

TNF-L or TNFR-L proteins or polypeptides can be isolated from TNF-L and TNFR-L-producing cells, such as spleen, thymus, prostate, colon, ovary, small intestine, peripheral blood lymphocytes, or from cell lines such as K562 (chronic myeloleukemia), G361 (melanoma), or SW480 (colorectal adenocarcinoma), using biochemical methods which are standard in the art. These methods include, but are not limited to, size exclusion chromatography, ammonium sulfate fractionation, ion exchange chromatography, affinity chromatography, crystallization, electrofocusing, and preparative gel electrophoresis. The skilled artisan can readily select methods which will result in a preparation of TNF-L or TNFR-L protein which is substantially free from other proteins and from carbohydrates, lipids, or subcellular organelles. A preparation of

isolated and purified TNF-L or TNFR-L protein is at least 80% pure; preferably, the preparations are 90%, 95%, or 99% pure. Purity of the preparations can be assessed by any means known in the art, such as SDS-polyacrylamide gel electrophoresis.

Human TNF-L and human or mammalian TNFR-L proteins, polypeptides, or
5 variants can be produced by recombinant DNA methods or by synthetic chemical methods. For production of recombinant TNF-L or TNFR-L proteins or polypeptides, coding sequences selected from the nucleotide sequences shown in SEQ ID NOS:6-10 can be expressed in known prokaryotic or eukaryotic expression systems. Bacterial, yeast, insect, or mammalian expression systems can be used, as is known in the art.

10 Alternatively, synthetic chemical methods, such as solid phase peptide synthesis, can be used to synthesize human TNF-L or human or mammalian TNFR-L protein, polypeptides, or variants. General means for the production of peptides, analogs or derivatives are outlined in CHEMISTRY AND BIOCHEMISTRY OF AMINO ACIDS, PEPTIDES, AND PROTEINS -- A SURVEY OF RECENT DEVELOPMENTS, B. Weinstein, ed. (1983).
15 Substitution of D-amino acids for the normal L-stereoisomer of a TNF-L or TNFR-L protein of the invention can be carried out to increase the half-life of the molecule.

Fusion proteins comprising at least 6, 7, 8, 9, 10, 12, 15, 20, 25, 30, 35, 40, 50, 75, 100, 125, 150, 175, 200, 225, 250, or 275 contiguous amino acids of SEQ ID NO:1, at least 6, 7, 8, 9, 10, 12, 15, 20, 25, 30, 35, 40, 50, 75, 100, 125, or 150 contiguous
20 amino acids of SEQ ID NO:2, at least 6, 7, 8, 9, 10, 12, 15, 20, 25, 30, 35, 40, 50, 75, 100, 125, 150, 175, or 200 contiguous amino acids of SEQ ID NO:3, at least 6, 7, 8, 9, 10, 12, 15, 20, 25, 30, 35, 40, 50, 75, 100, 125, 130, or 140 contiguous amino acids of SEQ ID NO:4, or at least 6, 7, 8, 9, 10, 12, 15, 20, 25, 30, 35, 40, 50, 75, 100, 125, 150, 175, or 200 contiguous amino acids of SEQ ID NO:5 can also be constructed. TNF-L
25 and TNFR-L fusion proteins are useful for generating antibodies against TNF-L and TNFR-L amino acid sequences and for use in various assay systems. For example, TNF-L and TNFR-L fusion proteins can be used to identify proteins which interact with these proteins which influence their biological activity and/or ability to bind to their respective binding partners. Physical methods, such as protein affinity chromatography, or library-
30 based assays for protein-protein interactions such as the yeast two-hybrid or phage

display systems, can also be used for this purpose. Such methods are well known in the art and can also be used as drug screens.

A TNF-L or TNFR-L fusion protein comprises two protein segments fused together by means of a peptide bond. The first protein segment consists of at least 6, 7, 8, 9, 10, 12, 15, 20, 25, 30, 35, 40, 50, 75, 100, 125, 150, 175, 200, 225, 250, or 275
5 contiguous amino acids of SEQ ID NO:1, at least 6, 7, 8, 9, 10, 12, 15, 20, 25, 30, 35, 40, 50, 75, 100, 125, or 150 contiguous amino acids of SEQ ID NO:2, at least 6, 7, 8, 9, 10, 12, 15, 20, 25, 30, 35, 40, 50, 75, 100, 125, 150, 175, or 200 contiguous amino acids of SEQ ID NO:3, at least 6, 7, 8, 9, 10, 12, 15, 20, 25, 30, 35, 40, 50, 75, 100, 125, 130, or
10 140 contiguous amino acids of SEQ ID NO:4, or at least 6, 7, 8, 9, 10, 12, 15, 20, 25, 30, 35, 40, 50, 75, 100, 125, 150, 175, or 200 contiguous amino acids of SEQ ID NO:5. The amino acids can be selected from the amino acid sequences shown in SEQ ID NOS:1-5 or from a biologically active variants of those sequences. The first protein segment can also be a full-length TNF-L or TNFR-L protein comprising an amino acid sequence as
15 shown in SEQ ID NOS:1-5. The first protein segment can be N-terminal or C-terminal, as is convenient.

The second protein segment can be a full-length protein or a protein fragment or polypeptide. Proteins commonly used in fusion protein construction include β -galactosidase, β -glucuronidase, green fluorescent protein (GFP), autofluorescent
20 proteins, including blue fluorescent protein (BFP), glutathione-S-transferase (GST), luciferase, horseradish peroxidase (HRP), and chloramphenicol acetyltransferase (CAT). Epitope tags can be used in fusion protein constructions, including histidine (His) tags, FLAG tags, influenza hemagglutinin (HA) tags, Myc tags, VSV-G tags, and thioredoxin (Trx) tags. Other fusion constructions can include maltose binding protein (MBP), S-tag,
25 Lex A DNA binding domain (DBD) fusions, GAL4 DNA binding domain fusions, and herpes simplex virus (HSV) BP16 protein fusions.

According to one particularly preferred embodiment, a TNFR-L protein is fused to the Fc domain of an IgG1 molecule. Such a fusion protein is be useful for inhibiting the action of a TNF ligand.

30 TNF-L or TNFR-L fusion proteins can be made by covalently linking the first

and second protein segments or by standard procedures in the art of molecular biology. Recombinant DNA methods can be used to prepare the fusion proteins, for example, by making a DNA construct which comprises coding sequences selected from SEQ ID NOS:6-10 in proper reading frame with nucleotides encoding the second protein segment
5 and expressing the DNA construct in a host cell, as is known in the art. Many kits for constructing fusion proteins are available from companies which supply research labs with tools for experiments, including, for example, Promega Corporation (Madison, WI), Stratagene (La Jolla, CA), Clontech (Mountain View, CA), Santa Cruz Biotechnology (Santa Cruz, CA), MBL International Corporation (MIC; Watertown, MA), and
10 Quantum Biotechnologies (Montreal, Canada; 1-888-DNA-KITS).

Isolated TNF-L or TNFR-L proteins, polypeptides, biologically active variants, or fusion proteins can be used as immunogens, to obtain a preparation of antibodies which specifically bind to epitopes of TNF-L or TNFR-L proteins. The antibodies can be used, *inter alia*, to detect TNF-L or TNFR-L proteins in tissue of humans or other
15 mammals or in fractions thereof. The antibodies can also be used to detect the presence of mutations in genes which result in under- or over-expression of TNF-L or TNFR-L proteins or in expression of a TNF-L or TNFR-L protein with altered size or electrophoretic mobility. By binding to TNF-L or TNFR-L proteins, antibodies can also alter the binding properties or biological functions of the proteins, for example for
20 therapeutic use.

Antibodies which specifically bind to epitopes of TNF-L or TNFR-L proteins, polypeptides, fusion proteins, or biologically active variants can be used in immunochemical assays, including but not limited to Western blots, ELISAs, radioimmunoassays, immunohistochemical assays, immunoprecipitations, or other
25 immunochemical assays known in the art. Typically, antibodies provide a detection signal at least 5-, 10-, or 20-fold higher than a detection signal provided with other proteins when used in such immunochemical assays. Preferably, antibodies which specifically bind to TNF-L or TNFR-L protein epitopes do not detect other proteins in immunochemical assays and can immunoprecipitate TNF-L or TNFR-L proteins or
30 polypeptides from solution.

TNF-L- or TNFR-L-specific antibodies specifically bind to epitopes present in a protein having the amino acid sequence shown in SEQ ID NOS:1-5 or to biologically active variants of those sequences. Typically, at least 6, 8, 10, or 12 contiguous amino acids are required to form an epitope. However, epitopes which involve non-contiguous amino acids may require more, *e.g.*, at least 15, 25, or 50 amino acids. Preferably, TNF-L or TNFR-L epitopes are not present in other proteins.

Protein epitopes which are particularly antigenic can be selected, for example, by routine screening of polypeptides for antigenicity or by applying a theoretical method for selecting antigenic regions of a protein to the amino acid sequences shown in SEQ ID NOS:1-5. Such methods are taught, for example, in Hopp and Wood, *Proc. Natl. Acad. Sci. U.S.A.* 78, 3824-28 (1981), Hopp and Wood, *Mol. Immunol.* 20, 483-89 (1983), and Sutcliffe *et al.*, *Science* 219, 660-66 (1983).

Any type of antibody known in the art can be generated to bind specifically to TNF-L or TNFR-L epitopes. For example, preparations of polyclonal and monoclonal antibodies can be made using standard methods which are well known in the art. Similarly, single-chain antibodies can also be prepared. Single-chain antibodies which specifically bind to TNF-L or TNFR-L epitopes can be isolated, for example, from a single-chain immunoglobulin display library, as is known in the art. The library is "panned" against the amino acid sequences disclosed herein, and a number of single chain antibodies which bind with high-affinity to different epitopes of proteins of the invention can be isolated. Hayashi *et al.*, 1995, *Gene* 160:129-30. Single-chain antibodies can also be constructed using a DNA amplification method, such as the polymerase chain reaction (PCR), using hybridoma cDNA as a template. Thirion *et al.*, 1996, *Eur. J. Cancer Prev.* 5:507-11.

Single-chain antibodies can be mono- or bispecific, and can be bivalent or tetravalent. Construction of tetravalent, bispecific single-chain antibodies is taught, for example, in Coloma and Morrison, 1997, *Nat. Biotechnol.* 15:159-63. Construction of bivalent, bispecific single-chain antibodies is taught *inter alia* in Mallender and Voss, 1994, *J. Biol. Chem.* 269:199-206.

A nucleotide sequence encoding a single-chain antibody can be constructed using

manual or automated nucleotide synthesis, cloned into an expression construct using standard recombinant DNA methods, and introduced into a cell to express the coding sequence, as described below. Alternatively, single-chain antibodies can be produced directly using, for example, filamentous phage technology. Verhaar *et al.*, 1995, *Int. J. Cancer* 61:497-501; Nicholls *et al.*, 1993, *J. Immunol. Meth.* 165:81-91.

Monoclonal and other antibodies can also be "humanized" in order to prevent a patient from mounting an immune response against the antibody when it is used therapeutically. Such antibodies may be sufficiently similar in sequence to human antibodies to be used directly in therapy or may require alteration of a few key residues. Sequence differences between, for example, rodent antibodies and human sequences can be minimized by replacing residues which differ from those in the human sequences, for example, by site directed mutagenesis of individual residues, or by grafting of entire complementarity determining regions. Alternatively, one can produce humanized antibodies using recombinant methods, as described in GB2188638B. Antibodies which specifically bind to TNF-L or TNFR-L epitopes can contain antigen binding sites which are either partially or fully humanized, as disclosed in U.S. 5,565,332.

Other types of antibodies can be constructed and used in methods of the invention. For example, chimeric antibodies can be constructed as disclosed, for example, in WO 93/03151. Binding proteins which are derived from immunoglobulins and which are multivalent and multispecific, such as the "diabodies" described in WO 94/13804, can also be prepared.

Antibodies can be purified by methods well known in the art. For example, antibodies can be affinity purified by passing the antibodies over a column to which a TNF-L or TNFR-L protein, polypeptide, biologically active variant, or fusion protein is bound. The bound antibodies can then be eluted from the column, using a buffer with a high salt concentration.

TNF-L- or TNFR-L-specific binding polypeptides other than antibodies can also be identified. These polypeptides include ligands of TNFR-L proteins and receptors of TNF-L proteins. TNF-L- or TNFR-L-specific binding polypeptides are polypeptides which bind with TNF-L or TNFR-L proteins or their variants and which have a

measurably higher binding affinity for TNF-L or TNFR-L and polypeptide variants of these proteins than for other polypeptides tested for binding. Higher affinity by a factor of 10 is preferred, more preferably a factor of 100. Such polypeptides can be found, for example, using the yeast two-hybrid system.

5 Nucleotide sequences which encode TNF-L or TNFR-L proteins are shown in SEQ ID NOS:6-10. Isolated and purified polynucleotides according to the invention can be single- or double-stranded, are subgenomic, and contain less than a whole chromosome. Preferably, the subgenomic polynucleotides are intron-free.

Isolated and purified subgenomic polynucleotides according to the invention can
10 comprise at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1000, 1100, or 1200 contiguous nucleotides of SEQ ID NO:6, at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 75, 100, 150, 200, 250, 300, 350, 400, or 450 contiguous nucleotides of SEQ ID NO:7, at least 10, 11, 12, 13, 14, 15, 16, 17, 18,
15 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 575, 600, 650, 700, 750, 800, or 850 contiguous nucleotides of SEQ ID NO:8, at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 575, or 600 contiguous nucleotides of SEQ ID NO:9, or at least 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25,
20 30, 35, 40, 45, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 600, 700, 800, 900, 1000, 1100, or 1200 contiguous nucleotides of SEQ ID NO:10 or can comprise SEQ ID NOS:6, 7, 8, 9, or 10. Such polynucleotides can be used, for example, as primers or probes or for expression of TNF-L or TNFR-L proteins or polypeptides.

The complements of the nucleotide sequences shown in SEQ ID NOS:6-10 are
25 contiguous nucleotide sequences which form Watson-Crick base pairs with a contiguous nucleotide sequence as shown in SEQ ID NOS:6-10. The complements of SEQ ID NOS:6-10 are polynucleotides of the invention and can be used, for example, to provide antisense oligonucleotides, primers, and probes.

Antisense oligonucleotides, primers, and probes of the invention can consist of at
30 least 11, 12, 15, 20, 25, 30, 50, or 100 contiguous nucleotides which are complementary

to the coding sequences shown in SEQ ID NOS:6-10. A complement of the entire coding sequence can also be used. Double-stranded subgenomic polynucleotides which comprise all or a portion of the nucleotide sequences shown in SEQ ID NOS:6-10, as well as polynucleotides which encode TNF-L- or TNFR-L-specific antibodies or
5 ribozymes, are also subgenomic polynucleotides according to the invention.

Degenerate nucleotide sequences encoding amino acid sequences of proteins or biologically active protein variants as well as homologous nucleotide sequences which are at least 65%, 75%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the nucleotide sequences shown in SEQ ID NOS:6-10, are also subgenomic polynucleotides
10 according to the invention and can be used in the methods disclosed herein. Percent sequence identity between a nucleotide sequence of SEQ ID NOS:6-10 and a putative homologous or degenerate nucleotide sequence is determined using computer programs which employ the Smith-Waterman algorithm, for example as implemented in the MPSRCH program (Oxford Molecular), using an affine gap search with the following
15 parameters: a gap open penalty of 12 and a gap extension penalty of 1.

Nucleotide sequences which hybridize to the coding sequences shown in SEQ ID NOS:6-10 or their complements with at most 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, or 35% basepair mismatches are also *TNF-L* or *TNFR-L* subgenomic polynucleotides. For example, using the following wash conditions--2X SSC (0.3 M sodium chloride, 0.03 M
20 sodium citrate, pH 7.0), 0.1% SDS, room temperature twice, 30 minutes each; then 2X SSC, 0.1% SDS, 50 °C once, 30 minutes; then 2X SSC, room temperature twice, 10 minutes each--homologous *TNF-L* or *TNFR-L* sequences can be identified which contain at most about 25-30% basepair mismatches with SEQ ID NOS:6-10 or their complements. More preferably, homologous nucleic acid strands contain 15-25%
25 basepair mismatches, even more preferably 5-15% basepair mismatches.

Species homologs of *TNF-L* or *TNFR-L* subgenomic polynucleotides of the invention can also be identified by making suitable probes or primers and screening cDNA expression libraries from other species, such as mice, monkeys, yeast, or bacteria. It is well known that the T_m of a double-stranded DNA decreases by 1-1.5 °C with every
30 1% decrease in homology (Bonner *et al.*, *J. Mol. Biol.* 81, 123 (1973). Homologous

TNF-L or *TNFR-L* human polynucleotides or *TNF-L* or *TNFR-L* polynucleotides of other species can therefore be identified, for example, by hybridizing a putative homologous *TNF-L* or *TNFR-L* polynucleotide with a polynucleotide having a nucleotide sequence of SEQ ID NO:6, 7, 8, 9, or 10, comparing the melting temperature of the test hybrid with the melting temperature of a hybrid comprising a polynucleotide having a nucleotide sequence of SEQ ID NOS:6, 7, 8, 9, or 10 and a polynucleotide which is perfectly complementary to SEQ ID NO:6, 7, 8, 9, or 10, and calculating the number of basepair mismatches within the test hybrid.

Nucleotide sequences which hybridize to the coding sequences shown in SEQ ID NOS:6-10 or their complements following stringent hybridization and/or wash conditions are also *TNF-L* or *TNFR-L* subgenomic polynucleotides. Stringent wash conditions are well known and understood in the art and are disclosed, for example, in Sambrook *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL, 2d ed., 1989, at pages 9.50-9.51.

Typically, for stringent hybridization conditions a combination of temperature and salt concentration should be chosen that is approximately 12-20 °C below the calculated T_m of the hybrid under study. The T_m of a hybrid between a sequence shown in SEQ ID NO:6, 7, 8, 9, or 10 and a polynucleotide sequence which is 65%, 75%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to SEQ ID NO:6, 7, 8, 9, or 10 can be calculated, for example, using the equation of Bolton and McCarthy, *Proc. Natl. Acad. Sci. U.S.A.* 48, 1390 (1962):

$$T_m = 81.5\text{ }^{\circ}\text{C} - 16.6(\log_{10}[\text{Na}^+]) + 0.41(\%G + C) - 0.63(\%\text{formamide}) - 600/l, \quad \text{where } l = \text{the length of the hybrid in basepairs.}$$

Stringent wash conditions include, for example, 4X SSC at 65 °C, or 50% formamide, 4X SSC at 42 °C, or 0.5X SSC, 0.1% SDS at 65 °C. Highly stringent wash conditions include, for example, 0.2X SSC at 65 °C.

Subgenomic polynucleotides can be isolated and purified free from other nucleotide sequences using standard nucleic acid purification techniques. For example, restriction enzymes and probes can be used to isolate subgenomic polynucleotide fragments which comprise *TNF-L* or *TNFR-L* coding sequences. Isolated and purified

subgenomic polynucleotides are in preparations which are free or at least 90% free of other molecules.

Complementary DNA (cDNA) molecules which encode TNF-L or TNFR-L proteins are also *TNF-L* or *TNFR-L* subgenomic polynucleotides. cDNA molecules can be made with standard molecular biology techniques, using TNF-L or TNFR-L mRNA as a template. cDNA molecules can thereafter be replicated using molecular biology techniques known in the art and disclosed in manuals such as Sambrook *et al.*, 1989. An amplification technique, such as the polymerase chain reaction (PCR), can be used to obtain additional copies of subgenomic polynucleotides, using either human or mammalian genomic DNA or cDNA as a template.

Alternatively, synthetic chemistry techniques can be used to synthesize subgenomic polynucleotide molecules. The degeneracy of the genetic code allows alternate nucleotide sequences to be synthesized which will encode a TNF-L protein comprising an amino acid sequence shown in SEQ ID NOS:1, 2, 3, 4, or 5 or a biologically active variant of one of those sequences.

The invention also provides polynucleotide probes which can be used to detect TNF-L or TNFR-L sequences, for example, in hybridization protocols such as Northern or Southern blotting or *in situ* hybridizations. *TNF-L* or *TNFR-L* polynucleotide probes of the invention comprise at least 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, or 40 or more contiguous nucleotides selected from SEQ ID NOS:6-10. Polynucleotide probes can comprise a detectable label, such as a radioisotopic, fluorescent, enzymatic, or chemiluminescent label.

TNF-L or *TNFR-L* subgenomic polynucleotides can be propagated in vectors and cell lines using techniques well known in the art. Expression systems in bacteria include those described in Chang *et al.*, *Nature* (1978) 275: 615, Goeddel *et al.*, *Nature* (1979) 281: 544, Goeddel *et al.*, *Nucleic Acids Res.* (1980) 8: 4057, EP 36,776, U.S. 4,551,433, deBoer *et al.*, *Proc. Natl. Acad. Sci. USA* (1983) 80: 21-25, and Siebenlist *et al.*, *Cell* (1980) 20: 269.

Expression systems in yeast include those described in Hinnen *et al.*, *Proc. Natl. Acad. Sci. USA* (1978) 75: 1929; Ito *et al.*, *J. Bacteriol.* (1983) 153: 163; Kurtz *et al.*,

- Mol. Cell. Biol.* (1986) 6: 142; Kunze *et al.*, *J. Basic Microbiol.* (1985) 25: 141; Gleeson *et al.*, *J. Gen. Microbiol.* (1986) 132: 3459, Roggenkamp *et al.*, *Mol. Gen. Genet.* (1986) 202 :302) Das *et al.*, *J. Bacteriol.* (1984) 158: 1165; De Louvencourt *et al.*, *J. Bacteriol.* (1983) 154: 737, Van den Berg *et al.*, *Bio/Technology* (1990) 8: 135; Kunze *et al.*, *J. Basic Microbiol.* (1985) 25: 141; Cregg *et al.*, *Mol. Cell. Biol.* (1985) 5: 3376, U.S. 4,837,148, US 4,929,555; Beach and Nurse, *Nature* (1981) 300: 706; Davidow *et al.*, *Curr. Genet.* (1985) 10: 380, Gaillardin *et al.*, *Curr. Genet.* (1985) 10: 49, Ballance *et al.*, *Biochem. Biophys. Res. Commun.* (1983) 112: 284-289; Tilburn *et al.*, *Gene* (1983) 26: 205-221, Yelton *et al.*, *Proc. Natl. Acad. Sci. USA* (1984) 81: 1470-1474, Kelly and Hynes, *EMBO J.* (1985) 4: 475479; EP 244,234, and WO 91/00357.

- Expression of *TNF-L* or *TNFR-L* subgenomic polynucleotides in insects can be accomplished as described in U.S. 4,745,051, Friesen *et al.* (1986) "The Regulation of Baculovirus Gene Expression" in: THE MOLECULAR BIOLOGY OF BACULOVIRUSES (W. Doerfler, ed.), EP 127,839, EP 155,476, and Vlak *et al.*, *J. Gen. Virol.* (1988) 69: 765-776, Miller *et al.*, *Ann. Rev. Microbiol.* (1988) 42: 177, Carbonell *et al.*, *Gene* (1988) 73: 409, Maeda *et al.*, *Nature* (1985) 315: 592-594, Lebacqz-Verheyden *et al.*, *Mol. Cell. Biol.* (1988) 8: 3129; Smith *et al.*, *Proc. Natl. Acad. Sci. USA* (1985) 82: 8404, Miyajima *et al.*, *Gene* (1987) 58: 273; and Martin *et al.*, *DNA* (1988) 7:99. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts are described in Luckow *et al.*, *Bio/Technology* (1988) 6: 47-55, Miller *et al.*, in GENETIC ENGINEERING (Setlow, J.K. *et al.* eds.), Vol. 8 (Plenum Publishing, 1986), pp. 277-279, and Maeda *et al.*, *Nature*, (1985) 315: 592-594.

- Mammalian expression of *TNF-L* or *TNFR-L* subgenomic polynucleotides can be accomplished as described in Dijkema *et al.*, *EMBO J.* (1985) 4: 761, Gorman *et al.*, *Proc. Natl. Acad. Sci. USA* (1982b) 79: 6777, Boshart *et al.*, *Cell* (1985) 41: 521 and U.S. 4,399,216. Other features of mammalian expression can be facilitated as described in Ham and Wallace, *Meth. Enz.* (1979) 58: 44, Barnes and Sato, *Anal. Biochem.* (1980) 102: 255, U.S. 4,767,704, US 4,657,866, US 4,927,762, US 4,560,655, WO 90/103430, WO 87/00195, and U.S. RE 30,985.

- TNF-L* or *TNFR-L* subgenomic polynucleotides can be on linear or circular

molecules. They can be on autonomously replicating molecules or on molecules without replication sequences. They can be regulated by their own or by other regulatory sequences, as is known in the art. *TNF-L* or *TNFR-L* subgenomic polynucleotides can be introduced into suitable host cells using a variety of techniques which are available in the art, such as transferrin-polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated DNA transfer, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, and calcium phosphate-mediated transfection.

Polynucleotides of the invention can also be used in gene delivery vehicles, for the purpose of delivering an mRNA or oligonucleotide (either with the sequence of a native mRNA or its complement), full-length protein, fusion protein, polypeptide, or ribozyme, or single-chain antibody, into a cell, preferably a eukaryotic cell. According to the present invention, a gene delivery vehicle can be, for example, naked plasmid DNA, a viral expression vector comprising a polynucleotide of the invention, or a polynucleotide of the invention in conjunction with a liposome or a condensing agent.

In one embodiment of the invention, the gene delivery vehicle comprises a promoter and one of the polynucleotides disclosed herein. Preferred promoters are tissue-specific promoters and promoters which are activated by cellular proliferation, such as the thymidine kinase and thymidylate synthase promoters. Other preferred promoters include promoters which are activatable by infection with a virus, such as the α - and β -interferon promoters, and promoters which are activatable by a hormone, such as estrogen. Other promoters which can be used include the Moloney virus LTR, the CMV promoter, and the mouse albumin promoter.

A gene delivery vehicle can comprise viral sequences such as a viral origin of replication or packaging signal. These viral sequences can be selected from viruses such as astrovirus, coronavirus, orthomyxovirus, papovavirus, paramyxovirus, parvovirus, picornavirus, poxvirus, retrovirus, togavirus or adenovirus. In a preferred embodiment, the gene delivery vehicle is a recombinant retroviral vector. Recombinant retroviruses and various uses thereof have been described in numerous references including, for example, Mann *et al.*, *Cell* 33:153, 1983, Cane and Mulligan, *Proc. Nat'l. Acad. Sci.*

USA 81:6349, 1984, Miller *et al.*, *Human Gene Therapy* 1:5-14, 1990, U.S. Patent Nos. 4,405,712, 4,861,719, and 4,980,289, and PCT Application Nos. WO 89/02,468, WO 89/05,349, and WO 90/02,806. Numerous retroviral gene delivery vehicles can be utilized in the present invention, including for example those described in EP 0,415,731; 5 WO 90/07936; WO 94/03622; WO 93/25698; WO 93/25234; U.S. Patent No. 5,219,740; WO 9311230; WO 9310218; Vile and Hart, *Cancer Res.* 53:3860-3864, 1993; Vile and Hart, *Cancer Res.* 53:962-967, 1993; Ram *et al.*, *Cancer Res.* 53:83-88, 1993; Takamiya *et al.*, *J. Neurosci. Res.* 33:493-503, 1992; Baba *et al.*, *J. Neurosurg.* 79:729-735, 1993 (U.S. Patent No. 4,777,127, GB 2,200,651, EP 0,345,242 and WO91/02805).

10 Particularly preferred retroviruses are derived from retroviruses which include avian leukosis virus (ATCC Nos. VR-535 and VR-247), bovine leukemia virus (VR-1315), murine leukemia virus (MLV), mink-cell focus-inducing virus (Koch *et al.*, *J. Vir.* 49:828, 1984; and Oliff *et al.*, *J. Vir.* 48:542, 1983), murine sarcoma virus (ATCC Nos. VR-844, 45010 and 45016), reticuloendotheliosis virus (ATCC Nos VR-994, VR-770 15 and 45011), Rous sarcoma virus, Mason-Pfizer monkey virus, baboon endogenous virus, endogenous feline retrovirus (*e.g.*, RD114), and mouse or rat gL30 sequences used as a retroviral vector. Particularly preferred strains of MLV from which recombinant retroviruses can be generated include 4070A and 1504A (Hartley and Rowe, *J. Vir.* 19:19, 1976), Abelson (ATCC No. VR-999), Friend (ATCC No. VR-245), Graffi (Ru *et al.*, *J. Vir.* 67:4722, 1993; and Yantchev *Neoplasma* 26:397, 1979), Gross (ATCC No. 20 VR-590), Kirsten (Albino *et al.*, *J. Exp. Med.* 164:1710, 1986), Harvey sarcoma virus (Manly *et al.*, *J. Vir.* 62:3540, 1988; and Albino *et al.*, *J. Exp. Med.* 164:1710, 1986) and Rauscher (ATCC No. VR-998), and Moloney MLV (ATCC No. VR-190). A particularly preferred non-mouse retrovirus is Rous sarcoma virus. Preferred Rous 25 sarcoma viruses include Bratislava (Manly *et al.*, *J. Vir.* 62:3540, 1988; and Albino *et al.*, *J. Exp. Med.* 164:1710, 1986), Bryan high titer (*e.g.*, ATCC Nos. VR-334, VR-657, VR-726, VR-659, and VR-728), Bryan standard (ATCC No. VR-140), Carr-Zilber (Adighitov *et al.*, *Neoplasma* 27:159, 1980), Engelbreth-Holm (Laurent *et al.*, *Biochem Biophys Acta* 908:241, 1987), Harris, Prague (*e.g.*, ATCC Nos. VR-772, and 45033), and 30 Schmidt-Ruppin (*e.g.* ATCC Nos. VR-724, VR-725, VR-354) viruses.

Any of the above retroviruses can be readily utilized in order to assemble or construct retroviral gene delivery vehicles given the disclosure provided herein and standard recombinant techniques (*e.g.*, Sambrook *et al.*, 1989, and Kunkle, *Proc. Natl. Acad. Sci. U.S.A.* 82:488, 1985) known in the art. Portions of retroviral expression
5 vectors can be derived from different retroviruses. For example, retrovector LTRs can be derived from a murine sarcoma virus, a tRNA binding site from a Rous sarcoma virus, a packaging signal from a murine leukemia virus, and an origin of second strand synthesis from an avian leukosis virus. These recombinant retroviral vectors can be used to generate transduction competent retroviral vector particles by introducing them into
10 appropriate packaging cell lines (*see* Serial No. 07/800,921, filed November 29, 1991). Recombinant retroviruses can be produced which direct the site-specific integration of the recombinant retroviral genome into specific regions of the host cell DNA. Such site-specific integration can be mediated by a chimeric integrase incorporated into the retroviral particle (*see* Serial No. 08/445,466 filed May 22, 1995). It is preferable that
15 the recombinant viral gene delivery vehicle is a replication-defective recombinant virus.

Packaging cell lines suitable for use with the above-described retroviral gene delivery vehicles can be readily prepared (*see* Serial No. 08/240,030, filed May 9, 1994; *see also* WO 92/05266) and used to create producer cell lines (also termed vector cell lines or "VCLs") for production of recombinant viral particles. In particularly preferred
20 embodiments of the present invention, packaging cell lines are made from human (*e.g.*, HT1080 cells) or mink parent cell lines, thereby allowing production of recombinant retroviral gene delivery vehicles which are capable of surviving inactivation in human serum. The construction of recombinant retroviral gene delivery vehicles is described in detail in WO 91/02805. These recombinant retroviral gene delivery vehicles can be used
25 to generate transduction competent retroviral particles by introducing them into appropriate packaging cell lines (*see* Serial No. 07/800,921). Similarly, adenovirus gene delivery vehicles can also be readily prepared and utilized given the disclosure provided herein (*see also* Berkner, *Biotechniques* 6:616-627, 1988, and Rosenfeld *et al.*, *Science* 252:431-434, 1991, WO 93/07283, WO 93/06223, and WO 93/07282).

30 A gene delivery vehicle can also be a recombinant adenoviral gene delivery

vehicle. Such vehicles can be readily prepared and utilized given the disclosure provided herein (see Berkner, *Biotechniques* 6:616, 1988, and Rosenfeld *et al.*, *Science* 252:431, 1991, WO 93/07283, WO 93/06223, and WO 93/07282). Adeno-associated viral gene delivery vehicles can also be constructed and used to deliver proteins or polynucleotides of the invention to cells *in vitro* or *in vivo*. The use of adeno-associated viral gene delivery vehicles *in vitro* is described in Chatterjee *et al.*, *Science* 258: 1485-1488 (1992), Walsh *et al.*, *Proc. Nat'l. Acad. Sci.* 89: 7257-7261 (1992), Walsh *et al.*, *J. Clin. Invest.* 94: 1440-1448 (1994), Flotte *et al.*, *J. Biol. Chem.* 268: 3781-3790 (1993), Ponnazhagan *et al.*, *J. Exp. Med.* 179: 733-738 (1994), Miller *et al.*, *Proc. Nat'l Acad. Sci.* 91: 10183-10187 (1994), Einerhand *et al.*, *Gene Ther.* 2: 336-343 (1995), Luo *et al.*, *Exp. Hematol.* 23: 1261-1267 (1995), and Zhou *et al.*, *Gene Therapy* 3: 223-229 (1996). *In vivo* use of these vehicles is described in Flotte *et al.*, *Proc. Nat'l Acad. Sci.* 90: 10613-10617 (1993), and Kaplitt *et al.*, *Nature Genet.* 8:148-153 (1994).

In another embodiment of the invention, a gene delivery vehicle is derived from a togavirus. Preferred togaviruses include alphaviruses, in particular those described in U.S. Serial No. 08/405,627, filed March 15, 1995, WO 95/07994. Alpha viruses, including Sindbis and ELVS viruses can be gene delivery vehicles for polynucleotides of the invention. Alpha viruses are described in WO 94/21792, WO 92/10578 and WO 95/07994. Several different alphavirus gene delivery vehicle systems can be constructed and used to deliver polynucleotides to a cell according to the present invention. Representative examples of such systems include those described in U.S. Patents 5,091,309 and 5,217,879. Particularly preferred alphavirus gene delivery vehicles for use in the present invention include those which are described in WO 95/07994, and U.S. Serial No. 08/405,627.

Preferably, the recombinant viral vehicle is a recombinant alphavirus viral vehicle based on a Sindbis virus. Sindbis constructs, as well as numerous similar constructs, can be readily prepared essentially as described in U.S. Serial No. 08/198,450. Sindbis viral gene delivery vehicles typically comprise a 5' sequence capable of initiating Sindbis virus transcription, a nucleotide sequence encoding Sindbis non-structural proteins, a viral junction region inactivated so as to prevent fragment transcription, and a Sindbis

RNA polymerase recognition sequence. Optionally, the viral junction region can be modified so that polynucleotide transcription is reduced, increased, or maintained. As will be appreciated by those in the art, corresponding regions from other alphaviruses can be used in place of those described above.

5 The viral junction region of an alphavirus-derived gene delivery vehicle can comprise a first viral junction region which has been inactivated in order to prevent transcription of the polynucleotide and a second viral junction region which has been modified such that polynucleotide transcription is reduced. An alphavirus-derived vehicle can also include a 5' promoter capable of initiating synthesis of viral RNA from
10 cDNA and a 3' sequence which controls transcription termination.

Other recombinant togaviral gene delivery vehicles which can be utilized in the present invention include those derived from Semliki Forest virus (ATCC VR-67; ATCC VR-1247), Middleberg virus (ATCC VR-370), Ross River virus (ATCC VR-373; ATCC VR-1246), Venezuelan equine encephalitis virus (ATCC VR923; ATCC VR-1250;
15 ATCC VR-1249; ATCC VR-532), and those described in U.S. Patents 5,091,309 and 5,217,879 and in WO 92/10578. The Sindbis vehicles described above, as well as numerous similar constructs, can be readily prepared essentially as described in U.S. Serial No. 08/198,450.

Other viral gene delivery vehicles suitable for use in the present invention
20 include, for example, those derived from poliovirus (Evans *et al.*, *Nature* 339:385, 1989, and Sabin *et al.*, *J. Biol. Standardization* 1:115, 1973) (ATCC VR-58); rhinovirus (Arnold *et al.*, *J. Cell. Biochem.* L401, 1990) (ATCC VR-1110); pox viruses, such as canary pox virus or vaccinia virus (Fisher-Hoch *et al.*, *PROC. NATL. ACAD. SCI. U.S.A.* 86:317, 1989; Flexner *et al.*, *Ann. N.Y. Acad. Sci.* 569:86, 1989; Flexner *et al.*, *Vaccine*
25 8:17, 1990; U.S. 4,603,112 and U.S. 4,769,330; WO 89/01973) (ATCC VR-111; ATCC VR-2010); SV40 (Mulligan *et al.*, *Nature* 277:108, 1979) (ATCC VR-305), (Madzak *et al.*, *J. Gen. Vir.* 73:1533, 1992); influenza virus (Luytjes *et al.*, *Cell* 59:1107, 1989; McMichael *et al.*, *The New England Journal of Medicine* 309:13, 1983; and Yap *et al.*, *Nature* 273:238, 1978) (ATCC VR-797); parvovirus such as adeno-associated virus
30 (Samulski *et al.*, *J. Vir.* 63:3822, 1989, and Mendelson *et al.*, *Virology* 166:154, 1988)

(ATCC VR-645); herpes simplex virus (Kit *et al.*, *Adv. Exp. Med. Biol.* 215:219, 1989) (ATCC VR-977; ATCC VR-260); *Nature* 277: 108, 1979); human immunodeficiency virus (EPO 386,882, Buchschacher *et al.*, *J. Vir.* 66:2731, 1992); measles virus (EPO 440,219) (ATCC VR-24); A (ATCC VR-67; ATCC VR-1247), Aura (ATCC VR-368),
5 Bebaru virus (ATCC VR-600; ATCC VR-1240), Cabassou (ATCC VR-922), Chikungunya virus (ATCC VR-64; ATCC VR-1241), Fort Morgan (ATCC VR-924), Getah virus (ATCC VR-369; ATCC VR-1243), Kyzylagach (ATCC VR-927), Mayaro (ATCC VR-66), Mucambo virus (ATCC VR-580; ATCC VR-1244), Ndumu (ATCC VR-371), Pixuna virus (ATCC VR-372; ATCC VR-1245), Tonate (ATCC VR-925),
10 Trinité (ATCC VR-469), Una (ATCC VR-374), Whataroa (ATCC VR-926), Y-62-33 (ATCC VR-375), O'Nyong virus, Eastern encephalitis virus (ATCC VR-65; ATCC VR-1242), Western encephalitis virus (ATCC VR-70; ATCC VR-1251; ATCC VR-622; ATCC VR-1252), and coronavirus (Hamre *et al.*, *Proc. Soc. Exp. Biol. Med.* 121:190, 1966) (ATCC VR-740).

15 A polynucleotide of the invention can also be combined with a condensing agent to form a gene delivery vehicle. In a preferred embodiment, the condensing agent is a polycation, such as polylysine, polyarginine, polyornithine, protamine, spermine, spermidine, and putrescine. Many suitable methods for making such linkages are known in the art (see, for example, Serial No. 08/366,787, filed December 30, 1994).

20 In an alternative embodiment, a polynucleotide is associated with a liposome to form a gene delivery vehicle. Liposomes are small, lipid vesicles comprised of an aqueous compartment enclosed by a lipid bilayer, typically spherical or slightly elongated structures several hundred Angstroms in diameter. Under appropriate conditions, a liposome can fuse with the plasma membrane of a cell or with the
25 membrane of an endocytic vesicle within a cell which has internalized the liposome, thereby releasing its contents into the cytoplasm. Prior to interaction with the surface of a cell, however, the liposome membrane acts as a relatively impermeable barrier which sequesters and protects its contents, for example, from degradative enzymes. Additionally, because a liposome is a synthetic structure, specially designed liposomes
30 can be produced which incorporate desirable features. See Stryer, *Biochemistry*, pp. 236-

240, 1975 (W.H. Freeman, San Francisco, CA); Szoka *et al.*, *Biochim. Biophys. Acta* 600:1, 1980; Bayer *et al.*, *Biochim. Biophys. Acta* 550:464, 1979; Rivnay *et al.*, *Meth. Enzymol.* 149:119, 1987; Wang *et al.*, *PROC. NATL. ACAD. SCI. U.S.A.* 84: 7851, 1987, Plant *et al.*, *Anal. Biochem.* 176:420, 1989, and U.S. Patent 4,762,915. Liposomes can
5 encapsulate a variety of nucleic acid molecules including DNA, RNA, plasmids, and expression constructs comprising polynucleotides such those disclosed in the present invention.

Liposomal preparations for use in the present invention include cationic (positively charged), anionic (negatively charged) and neutral preparations. Cationic
10 liposomes have been shown to mediate intracellular delivery of plasmid DNA (Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* 84:7413-7416, 1987), mRNA (Malone *et al.*, *Proc. Natl. Acad. Sci. USA* 86:6077-6081, 1989), and purified transcription factors (Debs *et al.*, *J. Biol. Chem.* 265:10189-10192, 1990), in functional form. Cationic liposomes are readily available. For example, N[1-2,3-dioleoyloxy)propyl]-N,N,N-triethylammonium
15 (DOTMA) liposomes are available under the trademark Lipofectin, from GIBCO BRL, Grand Island, NY. See also Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* 91: 5148-5152, 1994. Other commercially available liposomes include Transfectace (DDAB/DOPE) and DOTAP/DOPE (Boehringer). Other cationic liposomes can be prepared from readily available materials using techniques well known in the art. See, e.g., Szoka *et al.*, *Proc.*
20 *Natl. Acad. Sci. USA* 75:4194-4198, 1978; and WO 90/11092 for descriptions of the synthesis of DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane) liposomes.

Similarly, anionic and neutral liposomes are readily available, such as from Avanti Polar Lipids (Birmingham, AL), or can be easily prepared using readily available materials. Such materials include phosphatidyl choline, cholesterol, phosphatidyl
25 ethanolamine, dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), dioleoylphosphatidyl ethanolamine (DOPE), among others. These materials can also be mixed with the DOTMA and DOTAP starting materials in appropriate ratios. Methods for making liposomes using these materials are well known in the art.

The liposomes can comprise multilamellar vesicles (MLVs), small unilamellar
30 vesicles (SUVs), or large unilamellar vesicles (LUVs). The various liposome-nucleic

acid complexes are prepared using methods known in the art. See, e.g., Straubinger *et al.*, METHODS OF IMMUNOLOGY (1983), Vol. 101, pp. 512-527; Szoka *et al.*, *Proc. Natl. Acad. Sci. USA* 87:3410-3414, 1990; Papahadjopoulos *et al.*, *Biochim. Biophys. Acta* 394:483, 1975; Wilson *et al.*, *Cell* 17:77, 1979; Deamer and Bangham, *Biochim.*

5 *Biophys. Acta* 443:629, 1976; Ostro *et al.*, *Biochem. Biophys. Res. Commun.* 76:836, 1977; Fraley *et al.*, *Proc. Natl. Acad. Sci. USA* 76:3348, 1979; Enoch and Strittmatter, *Proc. Natl. Acad. Sci. USA* 76:145, 1979; Fraley *et al.*, *J. Biol. Chem.* 255:10431, 1980; Szoka and Papahadjopoulos, *Proc. Natl. Acad. Sci. USA* 75:145, 1979; and Schaefer-Ridder *et al.*, *Science* 215:166, 1982.

10 In addition, lipoproteins can be included with a polynucleotide of the invention for delivery to a cell. Examples of such lipoproteins include chylomicrons, HDL, IDL, LDL, and VLDL. Mutants, fragments, or fusions of these proteins can also be used. Modifications of naturally occurring lipoproteins can also be used, such as acetylated LDL. These lipoproteins can target the delivery of polynucleotides to cells expressing
15 lipoprotein receptors. Preferably, if lipoproteins are included with a polynucleotide, no other targeting ligand is included in the composition.

In another embodiment, naked polynucleotide molecules are used as gene delivery vehicles, as described in WO 90/11092 and U.S. Patent 5,580,859. Such gene delivery vehicles can be either DNA or RNA and, in certain embodiments, are linked to
20 killed adenovirus. Curiel *et al.*, *Hum. Gene. Ther.* 3:147-154, 1992. Other suitable vehicles include DNA-ligand (Wu *et al.*, *J. Biol. Chem.* 264:16985-16987, 1989), lipid-DNA combinations (Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* 84:7413-7417, 1989), liposomes (Wang *et al.*, *Proc. Natl. Acad. Sci.* 84:7851-7855, 1987) and microprojectiles (Williams *et al.*, *Proc. Natl. Acad. Sci.* 88:2726-2730, 1991).

25 One can increase the efficiency of naked polynucleotide uptake into cells by coating the polynucleotides onto biodegradable latex beads. This approach takes advantage of the observation that latex beads, when incubated with cells in culture, are efficiently transported and concentrated in the perinuclear region of the cells. The beads will then be transported into cells when injected into muscle. Polynucleotide-coated
30 latex beads will be efficiently transported into cells after endocytosis is initiated by the

latex beads and thus increase gene transfer and expression efficiency. This method can be improved further by treating the beads to increase their hydrophobicity, thereby facilitating the disruption of the endosome and release of polynucleotides into the cytoplasm.

5 The newly identified receptor proteins play regulatory roles in cell proliferation and/or differentiation. For example, they can induce the production of cytokines, immunoglobulins, and cell surface antigens. The receptors can also play a role in the negative regulation of osteoclastogenesis. Soluble TNFR-like receptors can be useful in the neutralization of TNF or TNF-like ligands for the treatment of rheumatoid arthritis
10 and Crohn's disease. Similarly, restoring normal apoptosis to a cell via these receptors can be used to treat viral diseases.

 A variety of diseases and conditions can be treated by modulating the activity of TNF-L or TNFR-L proteins of the invention. For example, TNFL proteins induce apoptosis of activated T cells, but rescue resting T cells from apoptosis. A TNF-L
15 protein can therefore be used to treat autoimmune diseases, such as myasthenia gravis, insulin-dependent diabetes mellitus, rheumatoid arthritis, multiple sclerosis, and systemic lupus erythematosus. TNF-L proteins also have tumor stimulating properties. Tumors can therefore be treated by inhibiting expression or activity of a TNF-L protein. Similarly, reducing expression of a TNFR-L protein or blocking its ligand binding site
20 can be used to treat tumors, whereas increasing expression of a TNFR-L protein can be used to treat autoimmune diseases such as those disclosed above.

 In one embodiment of the invention, expression of a *TNF-L* or *TNFR-L* gene is decreased using a ribozyme, an RNA molecule with catalytic activity. See, e.g., Cech, 1987, *Science* 236: 1532-1539; Cech, 1990, *Ann. Rev. Biochem.* 59:543-568; Cech,
25 1992, *Curr. Opin. Struct. Biol.* 2: 605-609; Couture and Stinchcomb, 1996, *Trends Genet.* 12: 510-515. Ribozymes can be used to inhibit gene function by cleaving an RNA sequence, as is known in the art (e.g., Haseloff *et al.*, U.S. 5,641,673).

 The coding sequences shown in SEQ ID NOS:6-10 can be used to generate ribozymes which will specifically bind to mRNA transcribed from a *TNF-L* or *TNFR-L*
30 gene. Methods of designing and constructing ribozymes which can cleave other RNA

molecules in trans in a highly sequence specific manner have been developed and described in the art (see Haseloff *et al.* (1988), *Nature* 334:585-591). For example, the cleavage activity of ribozymes can be targeted to specific RNAs by engineering a discrete "hybridization" region into the ribozyme. The hybridization region contains a sequence complementary to the target RNA and thus specifically hybridizes with the target (see, for example, Gerlach *et al.*, EP 321,201). Longer complementary sequences can be used to increase the affinity of the hybridization sequence for the target. The hybridizing and cleavage regions of the ribozyme can be integrally related; thus, upon hybridizing to the target RNA through the complementary regions, the catalytic region of the ribozyme can cleave the target.

TNF-L and *TNFR-L*-specific ribozymes can be introduced into cells, such as neoplastic cells, as part of a DNA construct, as is known in the art. The DNA construct can also include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and a transcriptional terminator signal, for controlling transcription of the ribozyme in the cells.

Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, or calcium phosphate precipitation, can be used to introduce the ribozyme-containing DNA construct into cells whose division it is desired to decrease, as described above. Alternatively, if it is desired that the cells stably retain the DNA construct, the DNA construct can be supplied on a plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art.

Expression of a *TNF-L* or *TNFR-L* gene can also be altered using an antisense oligonucleotide sequence. The antisense sequence is complementary to at least a portion of the coding sequence of a gene having a coding sequence shown in SEQ ID NO:6-10. Preferably, the antisense oligonucleotide sequence is at least six nucleotides in length, but can be about 8, 12, 15, 20, 25, 30, 35, 40, 45, or 50 nucleotides long. Longer sequences can also be used. *TNF-L* or *TNFR-L* antisense oligonucleotide molecules can be provided in a DNA construct and introduced into cells whose division is to be decreased, as described above.

Antisense oligonucleotides can be composed of deoxyribonucleotides,

ribonucleotides, or a combination of both. Oligonucleotides can be synthesized manually or by an automated synthesizer, by covalently linking the 5' end of one nucleotide with the 3' end of another nucleotide with non-phosphodiester internucleotide linkages such as alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamidate, carboxymethyl esters, carbonates, and phosphate triesters. See Brown, 1994, *Meth. Mol. Biol.* 20:1-8; Sonveaux, 1994, *Meth. Mol. Biol.* 26:1-72; Uhlmann *et al.*, 1990, *Chem. Rev.* 90:543-583.

Precise complementarity is not required for successful duplex formation between an antisense molecule and the complementary coding sequence of a *TNF-L* or *TNFR-L* gene. Antisense molecules which comprise, for example, 2, 3, 4, or 5 or more stretches of contiguous nucleotides which are precisely complementary to a *TNF-L* or *TNFR-L* coding sequence, each separated by a stretch of contiguous nucleotides which are not complementary to adjacent *TNF-L* or *TNFR-L* coding sequences, can provide targeting specificity for *TNF-L* or *TNFR-L* mRNA. Preferably, each stretch of contiguous nucleotides is at least 4, 5, 6, 7, or 8 or more nucleotides in length. Non-complementary intervening sequences are preferably 1, 2, 3, or 4 nucleotides in length. One skilled in the art can easily use the calculated melting point of an antisense-sense pair to determine the degree of mismatching which will be tolerated between a particular antisense oligonucleotide and a particular *TNF-L* or *TNFR-L* coding sequence.

TNF-L or *TNFR-L* antisense oligonucleotides can be modified without affecting their ability to hybridize to a *TNF-L* or *TNFR-L* coding sequence. These modifications can be internal or at one or both ends of the antisense molecule. For example, internucleoside phosphate linkages can be modified by adding cholesteryl or diamine moieties with varying numbers of carbon residues between the amino groups and terminal ribose. Modified bases and/or sugars, such as arabinose instead of ribose, or a 3', 5'-substituted oligonucleotide in which the 3' hydroxyl group or the 5' phosphate group are substituted, can also be employed in a modified antisense oligonucleotide. These modified oligonucleotides can be prepared by methods well known in the art. Agrawal *et al.*, 1992, *Trends Biotechnol.* 10:152-158; Uhlmann *et al.*, 1990, *Chem. Rev.*

90:543-584; Uhlmann *et al.*, 1987, *Tetrahedron. Lett.* 215:3539-3542.

Antibodies which specifically bind to a TNF-L and TNFR-L protein can also be used to alter effective levels of *TNF-L* or *TNFR-L* gene expression. TNF-L and TNFR-L-specific antibodies bind to TNF-L and TNFR-L proteins and prevent the proteins from functioning in the cell. Construction of such antibodies is disclosed above.

Expression of an endogenous *TNF-L* or *TNFR-L* gene in a cell can also be altered by introducing in frame with the endogenous *TNF-L* or *TNFR-L* gene a DNA construct comprising a *TNF-L* or *TNFR-L* targeting sequence, a regulatory sequence, an exon, and an unpaired splice donor site by homologous recombination, such that a homologously recombinant cell comprising the DNA construct is formed. The new transcription unit can be used to turn the *TNF-L* or *TNFR-L* gene on or off as desired. This method of affecting endogenous gene expression is taught in U.S. Patent No. 5,641,670.

The targeting sequence is a segment of at least 10, 12, 15, 20, or 50 contiguous nucleotides selected from a nucleotide sequence shown in SEQ ID NO:6-10. The transcription unit is located upstream of a coding sequence of the endogenous *TNF-L* or *TNFR-L* gene. The exogenous regulatory sequence directs transcription of the coding sequence of the *TNF-L* or *TNFR-L* gene.

Preferably, the mechanism used to decrease expression of the *TNF-L* or *TNFR-L* gene, whether ribozyme, antisense nucleotide sequence, or antibody, decreases expression of the gene by 50%, 60%, 70%, or 80%. Most preferably, expression of the gene is decreased by 90%, 95%, 99%, or 100%. The effectiveness of the mechanism chosen to alter expression of the gene can be assessed using methods well known in the art, such as hybridization of nucleotide probes to mRNA of the gene, quantitative RT-PCR, or detection of a TNF-L and TNFR-L protein using specific antibodies of the invention.

TNF-L and TNFR-L proteins or subgenomic polynucleotides can be used in therapeutic compositions for treating a variety of TNF-mediated disorders. Therapeutic compositions of the invention which comprise TNF-L protein or TNF-L protein encoding polynucleotides can be used, for example, to treat disorders in which abnormal numbers of T cells become activated. Activated T-lymphocytes are associated with

disease in graft versus host reactions (*e.g.*, bone marrow transplantation) and most forms of autoimmunity, including but not restricted to, multiple sclerosis, rheumatoid arthritis, lupus, and myasthenia gravis. T-lymphocyte-mediated primary diseases, such as juvenile diabetes, can also be treated using TNF-L protein or protein encoding polynucleotides.

5 TNF-L and TNFR-L therapeutic compositions of the invention can comprise a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers are well known to those in the art. Such carriers include, but are not limited to, large, slowly metabolized macromolecules, such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, and inactive virus
10 particles. Pharmaceutically acceptable salts can also be used in the composition, for example, mineral salts such as hydrochlorides, hydrobromides, phosphates, or sulfates, as well as the salts of organic acids such as acetates, propionates, malonates, or benzoates.

 TNF-L or TNFR-L therapeutic compositions can also contain liquids, such as water, saline, glycerol, and ethanol, as well as substances such as wetting agents,
15 emulsifying agents, or pH buffering agents. Liposomes, such as those described in U.S. 5,422,120, WO 95/13796, WO 91/14445, or EP 524,968 B1, can also be used as a carrier for a therapeutic TNF-L or TNFR-L composition.

 Typically, a therapeutic TNF-L or TNFR-L composition is prepared as an injectable, either as a liquid solution or suspension; however, solid forms suitable for
20 solution in, or suspension in, liquid vehicles prior to injection can also be prepared. A TNF-L or TNFR-L composition can also be formulated into an enteric coated tablet or gel capsule according to known methods in the art, such as those described in U.S. 4,853,230, EP 225,189, AU 9,224,296, and AU 9,230,801.

 Proliferative disorders, such as neoplasias, dysplasias, and hyperplasias, can be
25 treated by administration of a therapeutic TNF-L and TNFR-L composition which will inhibit TNF-L activity or expression. Neoplasias which can be treated with the therapeutic composition include, but are not limited to, melanomas, squamous cell carcinomas, adenocarcinomas, hepatocellular carcinomas, renal cell carcinomas, sarcomas, myosarcomas, non-small cell lung carcinomas, leukemias, lymphomas,
30 osteosarcomas, central nervous system tumors such as gliomas, astrocytomas,

oligodendrogliomas, and neuroblastomas, tumors of mixed origin, such as Wilms' tumor and teratocarcinomas, and metastatic tumors.

Proliferative disorders which can be treated with a therapeutic TNF-L composition include disorders such as anhydric hereditary ectodermal dysplasia, congenital alveolar dysplasia, epithelial dysplasia of the cervix, fibrous dysplasia of bone, and mammary dysplasia. Hyperplasias, for example, endometrial, adrenal, breast, prostate, or thyroid hyperplasias, or pseudoepitheliomatous hyperplasia of the skin can be treated with TNF-L or TNFR-L therapeutic compositions.

Even in disorders in which *TNF-L* or *TNFR-L* mutations are not implicated, decreasing expression of a *TNF-L* gene or a *TNFR-L* gene or decreasing a TNF-L or TNFR-L protein function can have a therapeutic application. In these disorders, decreasing *TNF-L* or *TNFR-L* expression or function can help to suppress tumors. Similarly, in tumors in which *TNF-L* or *TNFR-L* expression is not aberrant, effecting *TNF-L* or *TNFR-L* downregulation or decrease of TNF-L or TNFR-L activity can suppress metastases.

Administration of therapeutic compositions of the invention can include local or systemic administration, including injection, oral administration, particle gun, or catheterized administration, and topical administration. Various methods can be used to administer a therapeutic composition directly to a specific site in the body. For example, a small metastatic lesion can be located and a therapeutic composition injected several times in several different locations within the body of tumor. Alternatively, arteries which serve a tumor can be identified, and a therapeutic composition injected into such an artery, in order to deliver the composition directly into the tumor.

A tumor which has a necrotic center can be aspirated and the composition injected directly into the now empty center of the tumor. A therapeutic composition can be directly administered to the surface of a tumor, for example, by topical application of the composition. X-ray imaging can be used to assist in certain of the above delivery methods. Combination therapeutic agents, such as an anti-TNF-L neutralizing antibody and another therapeutic agent, can be administered simultaneously or sequentially.

Alternatively, a therapeutic composition can be introduced into human cells ex

vivo, and the cells then replaced into the human. Cells can be removed from a variety of locations including, for example, from a selected tumor or from an affected organ. In addition, the therapeutic composition can be inserted into non-tumorigenic cells, for example, dermal fibroblasts or peripheral blood leukocytes. If desired, particular
5 fractions of cells such as a T cell subset or stem cells can also be specifically removed from the blood (*see*, for example, PCT WO 91/16116). The removed cells can then be contacted with the therapeutic composition utilizing any of the above-described techniques, followed by the return of the cells to the human, preferably to or within the vicinity of a tumor. The above-described methods can additionally comprise the steps of
10 depleting fibroblasts or other non-contaminating tumor cells subsequent to removing tumor cells from a human, and/or the step of inactivating the cells, for example, by irradiation.

Receptor-mediated targeted delivery of therapeutic compositions containing *TNF-L* or *TNFR-L* subgenomic polynucleotides to specific tissues can also be used. Receptor-mediated DNA delivery techniques are described in, for example, Findeis *et al.* (1993),
15 *Trends in Biotechnol.* 11, 202-05; Chiou *et al.* (1994), GENE THERAPEUTICS: METHODS AND APPLICATIONS OF DIRECT GENE TRANSFER (J.A. Wolff, ed.); Wu & Wu (1988), *J. Biol. Chem.* 263, 621-24; Wu *et al.* (1994), *J. Biol. Chem.* 269, 542-46; Zenke *et al.* (1990), *Proc. Natl. Acad. Sci. U.S.A.* 87, 3655-59; Wu *et al.* (1991), *J. Biol. Chem.* 266,
20 338-42.

Both the dose of the *TNF-L* or *TNFR-L* composition and the means of administration can be determined based on the specific qualities of the therapeutic composition, the condition, age, and weight of the patient, the progression of the disease, and other relevant factors. If the composition contains *TNF-L* or *TNFR-L* protein,
25 polypeptide, or antibody, effective dosages of the composition are in the range of about 5 μg to about 50 $\mu\text{g/kg}$ of patient body weight, about 50 μg to about 5 mg/kg , about 100 μg to about 500 $\mu\text{g/kg}$ of patient body weight, and about 200 to about 250 $\mu\text{g/kg}$.

Therapeutic compositions containing *TNF-L* or *TNFR-L* subgenomic polynucleotides, ribozymes, or antisense oligonucleotides can be administered in a range
30 of about 100 ng to about 200 mg of DNA for local administration in a gene therapy

protocol. Concentration ranges of about 500 ng to about 50 mg, about 1 µg to about 2 mg, about 5 µg to about 500 µg, and about 20 µg to about 100 µg of DNA can also be used during a gene therapy protocol. Factors such as method of action and efficacy of transformation and expression are considerations that will effect the dosage required for ultimate efficacy of the *TNF-L* or *TNFR-L* subgenomic polynucleotides.

Where greater expression is desired over a larger area of tissue, larger amounts of a *TNF-L* or *TNFR-L* therapeutic composition or the same amount readministered in a successive protocol of administrations, or several administrations to different adjacent or close tissue portions of for example, a tumor site, may be required to effect a positive therapeutic outcome. In all cases, routine experimentation in clinical trials will determine specific ranges for optimal therapeutic effect.

The invention provides knock-out mammals whose endogenous *TNF-L* or *TNFR-L* gene is not expressed. Methods of making knock-out mammals are well known in the art. The mammal can be any experimental mammal, such as a mouse, rat, or rabbit; however, a mouse is preferred. The endogenous wild-type *TNF-L* or *TNFR-L* gene of the mammal can be deleted entirely, resulting in an absence of *TNF-L* or *TNFR-L* protein in the mammal. Alternatively, mutations such as deletions, insertions, missense substitutions, or inversions, can be introduced into a *TNF-L* or *TNFR-L* gene. Such mutations result in expression of truncated or otherwise aberrant forms of *TNF-L* or *TNFR-L* protein in the knock-out mammal. Mammalian cell lines which do not express an endogenous *TNF-L* or *TNFR-L* gene can also be constructed, as is known in the art.

Knock-out mammals and cells of the invention are useful as model systems for studying the effects of drugs in the absence of wild-type *TNF-L* or *TNFR-L* protein or in the presence of altered forms of the *TNF-L* or *TNFR-L* protein in the mammal or cell. Knock-out mammals can also be used to develop therapeutic treatments for diseases associated with alterations in *TNF-L* or *TNFR-L* gene expression, such as neoplasia or various autoimmune diseases.

The invention also provides screening methods which can be used to identify chemical agents which may have use in therapy. for example, regulators of the disclosed genes and proteins can be screened using a variety of methods. These

include ligand binding (Zhang *et al.*, *J. Biol. Chem.* 267:24069-24075), cytotoxicity (Creasey, *Cancer Res.* 47:145-149, 1987; Geigert, *Develop. Biol. Standard* 69:129; Tsujimoto, *J. Biochem.* 101:919-925, 1987; Kamijo, *Biochem. Biophys. Res. Commun.* 160:830-825, 1989; Sidhu, *Anticancer Res.* 9:1569-1576, 1989),
5 differentiation (Kamijo, 1989), maturation of osteoclasts from hematopoietic precursors (Lacey, *Endocrinology* 136:2367-2376, 1995), and proliferation (Tsujimoto, 1989).

The ability of a test compound or a potential therapeutic agent to stimulate or inhibit activity of a TNF-L or TNFR-L protein can be assessed by determining or
10 measuring the viability of the population of cells. A test compound which increases or decreases cell lysis or cell death is a modulator of the TNF-L or TNF-LR protein and can be used as a therapeutic agent to regulate TNF activities, such as cell lysis or cell death. A test compound which increases cell lysis or cell death may be particularly useful in treatment of neoplastic growth. The polypeptide of the invention can be
15 applied to the cell exogenously, or it can be expressed by a cell which has been transfected with a subgenomic polynucleotide encoding the polypeptide.

Methods for measuring the viability of cells can be any which are known in the art. Cell death can be determined by contacting the cell with a dye and viewing it under a microscope. Viable cells can be observed to have an intact membrane and do
20 not stain, whereas dying or dead cells having "leaky" membranes do stain. Incorporation of the dye by the cell indicates the death of the cell. The most common dye used in the art for this purpose is trypan blue. Viability of cells can also be determined by detecting DNA synthesis. Cells can be cultured in cell medium with labeled nucleotides, such as [³H]-thymidine. The uptake or incorporation of the
25 labeled nucleotides by cells indicates DNA synthesis and cell viability. Death of tumor cells *in vivo* can be monitored by observing regression or shrinkage of a tumor. Any suitable diagnostic technique can be applied.

Other cellular proteins which are involved in the same biological pathways can be identified by looking for proteins which interact with the disclosed polypeptides. Natural
30 ligands can therefore be identified for the receptor proteins, and natural receptor proteins

can be identified for the ligands. Complex formation can be detected *in vitro* or *in vivo*. Many methods for detecting formation of protein complexes are known in the art, and any such methods can be used. For example, the yeast two-hybrid system can be used in cells to detect proteins which interact with the disclosed ligands and receptors.

- 5 Alternatively, protein complex formation can be tested *in vitro* and complexes detected by altered mobility on non-denaturing gels, or by co-immunoprecipitation.

Expression of TNFR-L proteins can serve as a marker of neoplasia. TNFR-L proteins can be detected in body samples, including tissues, serum, urine, sputum, and feces, using immunological techniques. Expression can also be observed by measuring
10 or detecting mRNA encoding the receptors. Any suitable technique can be used including but not limited to Northern blotting and RT-PCR.

- A *TNF-L* or *TNFR-L* subgenomic polynucleotide can also be delivered to subjects for the purpose of screening test compounds for those which are useful for enhancing transfer of *TNF-L* or *TNFR-L* subgenomic polynucleotides to the cell or for enhancing
15 subsequent biological effects of *TNF-L* or *TNFR-L* subgenomic polynucleotides within the cell. Such biological effects include hybridization to complementary *TNF-L* or *TNFR-L* mRNA and inhibition of its translation, expression of a *TNF-L* or *TNFR-L* subgenomic polynucleotide to form a *TNF-L* or *TNFR-L* mRNA, single-chain antibody, ribozyme, oligonucleotide, or protein and/or replication and integration of a *TNF-L* or
20 *TNFR-L* subgenomic polynucleotide. The subject can be a cell culture or an animal, preferably a mammal, more preferably a human.

- Test compounds which can be screened include any substances, whether natural products or synthetic, which can be administered to the subject *in vitro* or *in vivo*. Libraries or mixtures of test compounds can be tested. The test compound can be a
25 pharmacologic agent already known in the art or can be a compound previously unknown to have any pharmacological activity. The test compound can be naturally occurring or designed in the laboratory. It can be isolated from microorganisms, animals, or plants, and can be produced recombinantly, or synthesized by chemical methods known in the art. Test compounds or substances can be delivered before, after, or concomitantly with
30 a *TNF-L* or *TNFR-L* subgenomic polynucleotide. They can be administered separately or

in admixture with a *TNF-L* or *TNFR-L* subgenomic polynucleotide.

Integration of a delivered *TNF-L* or *TNFR-L* subgenomic polynucleotide can be monitored by any means known in the art. For example, Southern blotting of the delivered *TNF-L* or *TNFR-L* subgenomic polynucleotide can be performed. A change in the size of the fragments of a delivered polynucleotide indicates integration. Replication of a delivered polynucleotide can be monitored *inter alia* by detecting incorporation of labeled nucleotides combined with hybridization to a *TNF-L* or *TNFR-L* probe. Expression of a *TNF-L* or *TNFR-L* subgenomic polynucleotide can be monitored by detecting production of *TNF-L* or *TNFR-L* mRNA which hybridizes to the delivered polynucleotide or by detecting *TNF-L* or *TNFR-L* protein. *TNF-L* or *TNFR-L* protein can be detected immunologically. Thus, the delivery of *TNF-L* or *TNFR-L* subgenomic polynucleotides according to the present invention provides an excellent system for screening test compounds for their ability to enhance transfer of *TNF-L* or *TNFR-L* polynucleotides to a cell, by enhancing delivery, integration, hybridization, expression, replication or integration in a cell *in vitro* or *in vivo* in an animal, preferably a mammal, more preferably a human.

The *TNFL1* gene (SEQ ID NO:6) maps to human chromosome 13q34. Polynucleotide probes of *TNFL1* can therefore be used to identify this region of chromosome 13 in metaphase spreads of human chromosomes. Preparations of human metaphase chromosomes can be prepared using standard cytogenetic techniques from human primary tissues or cell lines. Polynucleotide probes comprising at least 12 contiguous nucleotides selected from the nucleotide sequence shown in SEQ ID NO:6 are used to identify the human chromosome. The polynucleotide probes can be labeled, for example, with a radioactive, fluorescent, biotinylated, or chemiluminescent label, and detected by well known methods appropriate for the particular label selected. Protocols for hybridizing polynucleotide probes to preparations of metaphase chromosomes are also well known in the art. A polynucleotide probe will hybridize specifically to nucleotide sequences in the chromosome preparations which are complementary to the nucleotide sequence of the probe.

A polynucleotide probe which hybridizes specifically to human chromosome

region 13q34 hybridizes to nucleotide sequences present in the *TNFL1* gene and not to nucleotide sequences present in other human genes. A polynucleotide probe which hybridizes specifically to an *TNFL1* gene provides a detection signal at least 5-, 10-, or 20-fold higher than the background hybridization provided with non-*TNFL1* coding sequences.

A human chromosome which specifically hybridizes to an *TNFL1* polynucleotide probe is identified as a human chromosome 13. Preferably, the polynucleotide probe identifies the long arm of human chromosome 13. More preferably, the polynucleotide probe identifies a q34 region of human chromosome 13.

The complete contents of the references cited in this disclosure are expressly incorporated by reference herein. The following examples are illustrative and are not meant to limit the scope of the invention disclosed herein.

EXAMPLE 1

This example describes cloning of the full-length cDNA for *TNFL1*.

TNFL1 was first identified from a database of expressed sequence tags (ESTs) by its homology to other members of the TNF family. The full-length cDNA was isolated by screening a liver cDNA library applying the genetrapp technique (Gibco). A liver library from Gibco BRL was screened using the Genetrapp cDNA positive selection system (catalog no. 10356-020) and two oligonucleotide primers. The sequence of the biotinylated primer is: 5'AGGTCCATGTCTTTGGG3' (SEQ ID NO:11) the sequence of the non biotinylated primer is: 5'GGGGATGAATTGAGTCTG3' (SEQ ID NO:12). The product of the repair reaction was transformed, plated on LB+ Amp (100 µg/ml) plates. The colonies were analyzed by colony hybridization with a radioactive fragment prepared by PCR using the primers 5'GTGCCCTCGAAGAAAAAG3' (SEQ ID NO:13) and 5'GCAAGTTGGAGTTCATC3' (SEQ ID NO:14).

The longest open reading frame was 1280 bp long and contained a poly A tail as well as an in-frame stop codon at position -117 upstream of the ATG at position +1, suggesting that this clone was full-length. The nucleotide sequence surrounding this ATG also matched the Kozak consensus sequence. The open reading frame encodes a

protein of 285 amino acids which we named Tumor Necrosis Factor Like 1 (TNFL1) (Figure 1A).

The lack of a signal sequence at the N-terminus and the presence of an internal hydrophobic domain are indicative of a type II transmembrane structure, which is similar to the structure of most of the other members of the TNF family with the exception of lymphotoxin α . Two potential N-glycosylation sites were also identified in the extracellular region of the protein. When aligned with the extra-cellular domains from other members of the TNF family, the extracellular domain of the TNFL1 protein showed an overall homology of 28% to the proteolytically cleaved form of TNF (Figure 1B).

EXAMPLE 2

This example shows the tissue distribution of *TNFL1* mRNA.

Northern blots showing mRNAs from different tissues and cancerous cell lines were purchased from Clontech. A Northern blot with mRNAs from hematopoietic cell lines and various cell types of the immune system was prepared with 2 μ g of poly A mRNA.

A probe prepared by digestion of the TNFL1 cDNA with *Eco*RI and *Xho*I was labeled by random priming with 35 S and Klenow enzyme (Rediprime kit from Amersham). The hybridization was performed in the Expresshyb buffer purchased from Clontech.

A 3 kb messenger mRNA corresponding to *TNFL1* mRNA was detected mainly in the organs of the immune system, such as peripheral blood lymphocytes, spleen, and thymus, as well as in the small intestine and ovary (Figure 2A). Human *TNFL1* mRNA was also detected in a few human cancer cell lines such as the chronic myelogenous leukemia cell line K562 and the melanoma cell line G361 (Figure 2A). Mouse mRNA was detected in heart, spleen, and lung using as a probe a mouse EST sequence which is homologous to the human *TNFL1* sequence (Accession No. AA254417) (Figure 2B).

Because *TNFL1* mRNA was expressed in the spleen of both mouse and human samples as well as in peripheral blood leukocytes, a more precise analysis of the protein expression levels was carried out in the same tissues.

EXAMPLE 3

This example demonstrates expression of protein levels in mouse and human tissues.

5 A polyclonal antibody (D2710) was raised against amino acids 234-248 of TNFL1 and purified on a protein G column followed by a peptide affinity column. Amino acids 234-248 are highly conserved between the human and the mouse protein and differ by only 4 amino acids.

10 This antibody was able to recognize a purified 30 kDa TNFL1 protein by Western blot analysis (Figure 3A, lane 5). A single band corresponding to a 45 kDa protein was detected in cytoplasmic extracts from mouse bone marrow-derived dendritic cells and human monocytes. Both the 30 kDa and the 45 kDa bands were absent after incubation of D2710 with an excess of competitor peptide (Figure 3A, lanes 2, 4, and 6), suggesting that the 45 kDa protein corresponds to the full-length TNFL1 protein.

15 The affinity purified antibody was also able to detect TNFL1 expressed in insect cells. Insect cells were infected with a recombinant baculovirus expressing the TNFL1 protein and analyzed by flow cytometry. The protein TNFL1 was detected by intracellular staining with D2710 after fixation and permeabilization of the cells infected with the recombinant virus but not the wild type virus (Figure 3B).

20 Immunohistochemistry experiments were performed on sections from mouse spleen and lymphoid organs using the polyclonal antibody D2710. The spleen was isolated from an animal perfused with 4% paraformaldehyde in PBS, incubated in the same solution for one additional hour, and incubated overnight at 4 °C in a 10% sucrose solution. The spleen was then embedded in OCT prior to cryo-sectioning. The sections
25 were stored at -80 °C. Immunostaining was carried out using the following protocol. The sections were blocked in normal donkey IgG (whole molecule H+L; Jackson 017 000 003; lot 39113 at 25.5 g/l) diluted 1:100 in 1x PBS and Fc block diluted 1:50 (Pharmingen, catalog no. 01241A). The sections were then incubated with primary
30 antibodies D2710, anti-mouse CD11c (HL3, Pharmingen, catalog no. 09702D), anti-mouse Th1.2 CD90.2 (53-2.1, Pharmingen, catalog no. 01122A), or anti-mouse

CD45R/B220 (RA3-6B2, Pharmingen, catalog no. 01122A) diluted 1:50 in blocking reagent. The sections were washed three times for 3 minutes each in PBS.

The sections were then incubated in secondary antibody (biotin-labeled donkey anti-rabbit F(ab)₂ (Jackson 711-066-152) or biotin-labeled donkey anti-rat F(ab')₂ (Jackson 721-066-153) diluted 1:100 in PBS, washed 3 times for 3 min in PBS, and incubated for 30 minutes at room temperature with pre-equilibrated ABC-Alkaline Phosphatase reagent (Vector) or ABC-peroxidase (Vector). The sections were again washed 3 times for 3 minutes in PBS and incubated in a color developing reaction with levamisole (Vector; SK 5000) using a Vector black AP substrate kit (SK5200), Vector red AP substrate kit (SK 5100), or Vector AEC peroxidase substrate kit (SK4200). After washing again in PBS, alkaline phosphatase stained sections were counterstained with hematoxylin nuclear counterstain (Vector; H3401) and methyl green (Vector; H3402). Other sections were mounted in fluoromount (Southern Biotechnology Associates, catalog no. 100-01; Fisher OB100-01).

Normal rabbit IgG (R& D Systems, catalog no. AB-105C) at a 1:50 dilution, secondary antibody at a dilution of 1:100, and secondary antibody alone at a dilution of 1:100 were used as negative controls.

TNFL1 was constitutively and specifically expressed as a cell surface-bound protein in normal spleen (Figure 3C), but was weakly expressed in lymph nodes, mesenteric lymph nodes, and Peyer's patches. In the spleen, the pattern of expression was restricted to the marginal zone and the red pulp. The region stained with monoclonal antibodies directed against markers of the T cell population and of the B cell zone (Th1-2 and B220 respectively) did not overlap with the region stained with the antibody specific for TNFL1 (Figure 3D). A monoclonal antibody directed against the dendritic cell marker CD11c stained the T cell area and the marginal zone, as well as some isolated cell in the red pulp (Figure 3D).

Although it is possible that some cell sub-types present both the CD11c antigen and the TNFL1 protein at their surface, TNFL1 does not seem to be an exclusive marker of dendritic cells in the spleen. The Mac-3 antigen, a marker for macrophages and monocytes mainly localized in the red pulp, showed a pattern of expression very similar

to the one observed with TNFL1 (Figure 3D). Overall, these results suggest that TNFL1 is expressed at the surface of splenic macrophages, monocytes, or dendritic cells usually present in the marginal zone and the red pulp.

Flow cytometry experiments were performed on human PBMCs isolated from whole blood by Ficoll gradient centrifugation and the PBMCs analyzed. TNFL1 was found to be constitutively expressed on monocytes and B cells, but not on resting CD4+ and CD8+ T cells. Mouse bone marrow-derived dendritic cells cultivated for 10 days in the presence of GM-CSF also showed some surface staining with the TNFL1 antibody (Figure 4A). After incubation of PBMCs with anti-CD3 and anti-CD28 antibodies for 6 days in the presence of IL2, TNFL1 was shown to be upregulated at the surface of T cells (Figure 4B).

EXAMPLE 4

This example demonstrates expression of a soluble form of TNFL1 in the periplasm of *E. coli*.

A chimeric soluble version of TNFL1 was expressed in *E. coli* as a fusion protein comprising the extracellular portion of TNFL1 (amino acids 113-285; Figure 5A) and the *pelB* signal sequence for periplasmic localization. The EYMPMD peptide (SEQ ID NO:15) was inserted between the signal sequence and the TNFL1 sequence for convenient affinity purification. The cDNA for TNFL1 was cloned into the vector pET-22b(+) from Invitrogen, which contains the *pelB* signal sequence for periplasmic localization. A 100 ml culture was grown at 37 °C until it reached an OD of 0.7-0.9 and then grown at 25 °C for 24 hours after induction by 1 mM IPTG. The pellet was centrifuged at 4000 x g for 10 minutes and resuspended in 10 ml 30 mM Tris HCl, 20% sucrose, pH 8.0. After addition of 1 mM EDTA, the sample was incubated at room temperature for 5-10 min. The sample was then centrifuged at 8000 x g at 4 °C for 10 min. The supernatant was removed, and the pellet was resuspended in 10 ml ice-cold 5 mM MgSO₄. After a 10 minute incubation in an ice/water bath, the sample was centrifuged at 8000 x g at 4 °C for 10 min. The supernatant containing the periplasmic fraction was stored at -80 °C in 15 % glycerol. A control sample was processed in a

similar way with an empty vector construct.

A similar fusion construct was also designed for expression and purification from COS cells using the pSecTag vector (Invitrogen) with a signal sequence from the mouse Ig κ chain.

5

EXAMPLE 5

This example demonstrates purification of the TNFL1 fusion protein from *E. coli*.

BL21 (DE3) transformed *E. coli* were grown in a 10-l fermenter to an OD of 29-31 before induction with IPTG. The cells were harvested in a Beckman J-6B
10 centrifuge. The wet cell paste was subjected to osmotic shock treatment for periplasmic extraction (lot 10229-142) or lysozyme/EDTA spheroplast formation with retention of the spheroplast supernatant as the periplasmic fraction (lot 981001-M8). Using a Pall Filtron Centrasette tangential flow apparatus and 2 Centrasette 10 kD NMWCO membranes, the resulting periplasmic fraction was concentrated to 1 liter, then
15 buffer-exchanged by constant-flow diafiltration in the same apparatus against at least 6 volumes of PBS. The resulting solution was centrifuged at 10,000 rpm at 4 °C for 50 minutes in a Beckman J2-21 centrifuge with a JA-10 rotor.

The resulting supernatant was precipitated with 50% ammonium sulfate using an equal volume of saturated ammonium sulfate. The resulting pellet was resuspended in $\frac{1}{4}$
20 the original volume of PBS and loaded onto a glu-tag monoclonal antibody affinity column at a flow rate of 30 cm/hr. Following the load, a wash of 5-10 CV of PBS + 0.2% Tween 20 was performed, followed by 2 CV of PBS. Elution was effected by 5 CV of 0.1 mg/ml EYMPTD peptide (SEQ ID NO:16) (Research Genetics) followed by PBS. A strip of 1.5 CV 0.1 M glycine pH 2.7 was also collected into one-tenth volume
25 of 1 M Tris pH 8, and the column was neutralized with 1 CV of the same.

The peptide elution was concentrated in an Amicon 8400 stirred-cell with a YM-10 membrane. Peptide removal was effected by overnight dialysis of the concentrate against PBS (except for the samples treated to remove residual detergent). Endotoxin removal was by Triton X-114 phase separation. One-ninth volume of 10% Triton X-114
30 protein grade (Calbiochem) was added to samples chilled on ice. Constant agitation for

30 min. at 4 °C was followed by 15 min. incubation in a 37 °C water bath. Centrifugation at 10,000 x g recovered the supernatant as the detergent-poor aqueous fraction. The cycle was repeated 3-5 times, resulting in endotoxin levels of less than or equal to 0.05 EU/ml.

Detergent was removed by an anion-exchange chromatography step in which the protein was bound and eluted while the detergent and elution peptide flowed through. A Pharmacia HR 10/5 column packed with 1.7 ml TMAE-Fractogel (EM Merck) was equilibrated in 25 mM NaPi, pH 7.4, at 1 ml/minute on an Akta Explorer FPLC system. Fractions of 1 ml were collected across a 15 CV gradient from 0-0.6 M NaCl in 25 mM NaPi, pH 7.4. The protein eluted at a conductivity of about 22 mS/cm. The peak fractions were pooled and concentrated, if necessary, in a Amicon Centriplus-10 centrifugal concentration device.

The purified protein was detected in a Western blot with both the tag antibody and the D2710 antibody as a 30 kD protein (Figure 3A). A minor additional band at 21 kD was also observed. This band was identified by microsequencing as a degradation product of the 30 kD protein (Figure 5A).

EXAMPLE 6

This example demonstrates the effects of TNFL1 on T and B cells.

T cells isolated from healthy donors were activated for two days in the presence of anti-CD3 and anti-CD28 antibodies and incubated for an additional two days in the presence of different concentrations of purified TNFL1. Mouse and human T cells were purified by negative selection from PBMCs on a mouse or human T cell enrichment column (R&D Systems), and B cells were isolated by positive selection on Dynabeads M-450 Pan-B and detached from the beads with the polyclonal antibody Detachabead from Dynal. The T cells were activated for 2 days with anti-CD3 and anti-CD28 antibodies at a concentration of 10 µg/ml

T cell proliferation was assessed by thymidine incorporation at day 5. The assays were performed in a 96 well plate format with 100,000 cells/well. The thymidine incorporation assay was performed by addition of ³H-thymidine at a concentration of 1 µCi/well. After 8 hours, the cells were harvested onto a filter-paper and ³H-thymidine

uptake was measured by liquid scintillation counting.

The dose-response curve shows a 30-fold decrease in thymidine incorporation at the highest concentrations of TNFL1 (Figure 5B, left). Resting T cells incubated with the ligand for two days did not show any decrease in thymidine incorporation (Figure 5B, right).

In a separate experiment, the effect of TNFL1 was tested on anti-CD3- and anti-CD28-activated T cells and on anti-CD40-activated B cells. The T cells were activated for 2 days with anti-CD3 and anti-CD28 antibodies at a concentration of 10 µg/ml, and the B cells were activated for 2 days in the presence of anti-CD40 antibody at a concentration of 10 µg/ml. Both cell types showed a strong and comparable decrease in thymidine incorporation after addition of TNFL1 (Figure 5C). TNFL1 also inhibited the proliferation of murine T cells activated by allogeneic bone marrow-derived DCs by 90%.

We also tested the ability of TNFL1 to decrease the proliferation rate of mouse T cells activated by allogeneic bone marrow-derived dendritic cells in a mixed lymphocyte reaction (MLR). For the MLR reaction, 100,000 mouse T cells were mixed with 10,000 mouse dendritic cells. Dendritic cells were prepared from red blood cell-depleted bone marrow cells. The cells were resuspended in culture medium in a bacterial Petri dish at a concentration of 0.2×10^6 cells per ml. Recombinant mouse GM-CSF was added to the cells at a final concentration of 200 U/ml on days 1, 2, 3, 6 and 8. On day 10, LPS at 1 µg/ml or TNF-α at 500 U/ml was added to the cells. The cells were harvested on day 11.

In the absence of TNFL1, the cultured DCs triggered a 100-fold increase in thymidine incorporation in the responding T cells. When the assay was performed in the presence of TNFL1, the increase in thymidine incorporation was only 10-fold, showing that TNFL1 is able to reduce the T cell stimulation induced by allogeneic antigen presenting cells.

Overall, these results indicate that TNFL1 is able to induce a decrease in thymidine incorporation in activated B and T cells. This effect could be due to a direct inhibition of proliferation or to the induction of apoptosis of B and T cells.

In order to elucidate the exact mechanism of action of TNFL1 on activated T cells, a

TUNEL assay (TdT-mediated dUTP-FITC nick-end labeling; Boehringer Mannheim) was performed according to the manufacturer's specifications. Briefly, treated cells were washed twice in PBS supplemented with 1% bovine serum albumin, fixed with 0.4% paraformaldehyde, and permeabilized in 0.1% Triton X-100 in 0.1% sodium citrate.

5 TUNEL staining was performed with 50 μ l of TUNEL mix per sample at 37 °C for 1 hour. Negative controls were treated similarly but were not exposed to the enzyme.

After incubation of the T cells with TNFL1, the amount of apoptotic cells measured by flow cytometry analysis (Figure 5D). The blasted cells, visualized by forward size scattering (region R1), almost totally disappeared after addition of TNFL1.

10 At the same time, the amount of apoptotic, FITC-positive cells (region R2) dramatically increased. No apoptotic cells were detected in the absence of dUTP-FITC, whether or not TNFL1 was added.

These results strongly suggest that TNFL1 is able to induce apoptotic cell death of blasting as well as of slightly activated T cells.

15

EXAMPLE 7

This example demonstrates that TNFL1 activates NF κ B and leads to its translocation to the nucleus.

TNF activates the transcription factor NF κ B. NF κ B represents a family of

20 related proteins involved in the transcriptional control of numerous cellular genes such as interleukin-2, interleukin-2 receptor, γ -interferon, granulocyte macrophage colony-stimulating factor, histocompatibility antigens, TNF, and lymphotoxin

□□□□ We therefore decided to test whether TNFL1 could induce a similar intracellular response in Jurkat cells.

25 Jurkat cells (10^7) were incubated for one hour with PMA at a concentration of 1 μ g/ml or TNFL1 at a concentration of 3 μ g/ml in 1 ml of RPMI and 10% FBS. Nuclear extracts were prepared by centrifuging the cells at 2000 rpm and resuspending the pellet in 1 ml of cold buffer A (10 mM Hepes, 1.5 mM MgCl₂, 10 mM KCl, 0.5 mM DTT, 0.5 mM PMSF). After a second centrifugation at 2000 rpm, the pellet was resuspended in 20

30 μ l of buffer A and 0.1% NP40, incubated for 10 minutes on ice, and then centrifuged.

The pellet was resuspended in 15 μ l of buffer C (20 mM Hepes, pH 7.9, 0.42 M NaCl, 1.5 mM $MgCl_2$, 0.2 mM EDTA, pH 7.4, 25% glycerol, 0.5 mM PMSF, 0.5 mM DTT, 50 μ g/ml leupeptin, 50 μ g/ml pepstatin, and 78 μ g/ml benzamidin). The mixture was incubated for 15 minutes on ice, then centrifuged for 10 minutes at 14,000 rpm at 4 °C.

- 5 The supernatant was diluted in 75 μ l of buffer D (20 mM Hepes, pH 7.9, 20% glycerol, 0.2 mM EDTA, pH 7.4, 50 mM KCl, 0.5 mM DTT, and 0.5 mM PMSF).

- Five μ g of nuclear extract was incubated for 10 minutes at 4 °C in the presence of 5 μ g of polyIdC and wild-type or mutated competitor oligonucleotide (2 ng to 200 ng). Two-tenths of a nanogram of polynucleotide kinase-radiolabeled probe was added to the reaction, and the mixture was incubated at room temperature for 30 min.

- The mixture was loaded on a 5% acrylamide 60:1 x linked gel in Tris-glycine buffer, pH 8.3, and analyzed in an electrophoretic mobility shift assay. Specific bands, identified by competition with wild-type or mutated NF κ B binding sites, were observed with both PMA- and TNFL1-treated extracts (Figure 5E), indicating that TNFL1 is indeed able to activate NF κ B and leads to its translocation to the nucleus.

CLAIMS

1. An isolated human protein having an amino acid sequence which is at least 85% identical to an amino acid sequence selected from the group consisting of SEQ ID NOS:1 and 2, wherein percent identity is determined using a Smith-Waterman homology search algorithm using an affine gap search with a gap open penalty of 12 and a gap extension penalty of 1.
2. The isolated human protein of claim 1 which has an amino acid sequence selected from the group consisting of SEQ ID NOS:1 and 2.
3. A fusion protein comprising a first protein segment and a second protein segment fused together by means of a peptide bond, wherein the first protein segment consists of a protein having an amino acid sequence selected from the group consisting of SEQ ID NOS:1 and 2.
4. A preparation of antibodies which specifically bind to a protein having an amino acid sequence selected from the group consisting of SEQ ID NOS:1 and 2.
5. A cDNA molecule which encodes a protein having an amino acid sequence which is at least 85% identical to an amino acid sequence selected from the group consisting of SEQ ID NOS:1 and 2, wherein percent identity is determined using a Smith-Waterman homology search algorithm using an affine gap search with a gap open penalty of 12 and a gap extension penalty of 1.
6. The cDNA molecule of claim 5 which encodes an amino acid sequence selected from the group consisting of SEQ ID NOS:1 and 2.
7. The cDNA molecule of claim 6 which comprises a nucleotide sequence selected from the group consisting of SEQ ID NOS:6 and 7.
8. A cDNA molecule which is at least 85% identical to a nucleotide sequence selected from the group consisting of SEQ ID NOS:6 and 7, wherein percent identity is determined using a Smith-Waterman homology search algorithm using an affine gap search with a gap open penalty of 12 and a gap extension penalty of 1.
9. An isolated and purified subgenomic polynucleotide comprising a nucleotide

sequence which hybridizes to a nucleotide sequence selected from the group consisting of SEQ ID NOS:6 and 7 after washing with 0.2X SSC at 65 °C, wherein the nucleotide sequence encodes a protein having an amino acid sequence selected from the group consisting of SEQ ID NOS:1 and 2.

10. A construct comprising:

a promoter; and

a polynucleotide segment encoding an amino acid sequence selected from the group consisting of SEQ ID NOS:1 and 2, wherein the polynucleotide segment is located downstream from the promoter, wherein transcription of the polynucleotide segment initiates at the promoter.

11. A host cell comprising a construct which comprises:

a promoter; and

a polynucleotide segment encoding an amino acid sequence selected from the group consisting of SEQ ID NOS:1 and 2.

12. A recombinant host cell comprising a new transcription initiation unit, wherein the new transcription initiation unit comprises in 5' to 3' order:

(a) an exogenous regulatory sequence;

(b) an exogenous exon; and

(c) a splice donor site, wherein the new transcription initiation unit is located upstream of a coding sequence of a gene, wherein the coding sequence is selected from the group consisting of SEQ ID NOS:6 and 7, wherein the exogenous regulatory sequence controls transcription of the coding sequence of the gene.

13. A method of screening for a compound capable of modulating cell death inducing activity of a protein, comprising the steps of:

incubating a first population of cells and a protein in the presence of a test compound, wherein the protein comprises an amino acid sequence selected from the group of amino acid sequences shown in SEQ ID NOS:1-5;

incubating a second population of cells and the protein in the absence of a test compound; and

determining viability of the first and second populations, wherein a test compound which

increases or decreases viability of the first population relative to the second population is identified as capable of modulating the cell death inducing activity of the protein.

14. The method of claim 13 wherein the protein is provided to the first and second populations of cells by transfecting the first and second populations of cells with a polynucleotide encoding the protein.

15. The method of claim 13 wherein the polynucleotide comprises a nucleotide sequence selected from the group consisting of SEQ ID NOS:6-10.

16. A method of identifying a binding partner of a first protein, comprising the steps of:

incubating a first protein with a second protein, wherein the first protein comprises an amino acid sequence selected from the group consisting of SEQ ID NO: 1-5;
detecting formation of a complex between the first and second proteins, wherein formation of the complex identifies the second protein as a binding partner of the first protein.

A

1 MDDSTEREQSRLTSCLEKREEMKLEKCVSILPRKESPSVRSSKDGKLLAATLLALLSCC 60
 61 LTVVSFYQVAALOGDLASLRAELQGHAEKLPAGAGAPKAGLEEAPAVTAGLKIFEPPAP 120
 121 GEGNSSQNSRNKRAVQGPETVTQDCLQIADSETPTIQKGSYTFVPWLLSFKRGSAL EEKENKI 180
 181 KENKILVKETGYFFIYGQVLYTDKTYAMGHLIQKKVHVFGDELSLVTLCFRCIQNMPETL 240
 241 PNNSCYSAGIAKLEEGDELQLAIPRENAQISLDGDTVFFGALKLL 285


B

TNFI1) TVTQDCLQLIADSETPTIQKGSYTFVPWLLSFKRGSAL EEKENKI 185
 Fas ligand) KELRKVAHLTGKSNRSRSM--PLEWEDTYGIVL LSGVKYKKGG--L 181
 ltoxin alpha) STLKFAAHLIGDPSKQNS--L LWRANTDR AFLQDGFSL SNNS--L 99
 ltoxin beta) SPGLFAAHLIGAFLKQG--LGWETTKEQAF L TSGTQFSDAEG--L 125
 TNF) PSDKFAHVVANPQAEGQ--LQWLNRRANALLANGVE L RDNQ--L 126

TNFI1) L V K E T G Y F F I Y G Q V L Y T D - - - - - K T Y A M G H L I Q R K K V H V 219
 Fas ligand) V I N E T G L Y F V Y S K V Y F R G - - - - - Q S C N N L P L S H K V Y M R N S K Y 218
 ltoxin alpha) L V P T S G I Y F V Y S Q V V F S G K A Y - - S P K A T S S P L Y L A H E V O L F S S Q Y 163
 ltoxin beta) A L P Q D G L Y Y L Y C L V G Y R G R A P P G G D P Q G R S V T L R S S L Y R A G G A Y 170
 TNF) V V P S E G L Y L I Y S Q V L F R G Q G - - - - - C P S T H V L L T H T I S R I A V S Y 163

TNFI1) F G - D E L S L V T L F R - - - - - C I Q N M P E T L P N N S C Y S A G I A K L E E 255
 Fas ligand) P - - Q D L V M M E G K M - - - - - M S Y C T T G Q M W A R S S Y L G A V F N L T S 253
 ltoxin alpha) P - - F H V P L L S S Q K - - - - - M V Y P G L O E P W L H S M Y H G A A F O L T Q 177
 ltoxin beta) G P G T P E L L L E G A E T V T P V L D P A R R Q G Y G P L W Y T S V G F G G L V O L R R 215
 TNF) Q - - T K V N L L S A I K S P C Q R - - E T P E G A E A K P W Y E P I Y L G G V F O L E K 204

TNFI1) G D E L Q L A I P R E N A Q I S L D G D V T F F G A L K L L 285
 Fas ligand) A D H L Y V N V S E L S L V N F E E S - Q T F F G L Y N L 281
 ltoxin alpha) G D Q L S T H T D G I P H L V L S P S - T V F F G A F A L 205
 ltoxin beta) G E R V Y V N I S H P D M V D F A R C - K T F F G A V M V G 244
 TNF) G D R L S A E I N R P D Y L D F A E S G O V Y F G I I A L 233






Figure 3




Figure 5

SEQUENCE LISTING

SEQ ID NO: 1

MDDSTEREQSRLTSCCLKREEMKLKECVSILPRKESPSVRSSKDGKLLAATLLALLSCCLTVVSFYQVAA
 LOGDLASLRAELQGHHA EKLPAGAGAPKAGLEEAPAVTAGLKIFEPPAPGEGNSSQNSRNKRAVQGPEETV
 TQDCLQLIADSETPTIQKGSYTFVPWLLSFKRGSALEEKENKILVKETGYFFIYGQVLYTDKTYAMGHLIQ
 RKKVHVFGDELSLVTLFRCIQNMPETLPNNSCYSAGIAKLEEGDELQLAIPRENAQISLDGDVTFFGALKL
 L

SEQ ID NO:2:

Leu	Glu	Arg	Cys	Arg	Tyr	Cys	Asn	Val	Leu	Cys	Gly	Glu	Arg	Glu	Glu
1				5					10					15	
Glu	Ala	Arg	Ala	Cys	His	Ala	Thr	His	Asn	Arg	Ala	Cys	Arg	Cys	Arg
				20					25					30	
Thr	Gly	Phe	Phe	Ala	His	Ala	Gly	Phe	Cys	Leu	Glu	His	Ala	Ser	Cys
				35					40					45	
Pro	Pro	Gly	Ala	Gly	Val	Ile	Ala	Pro	Gly	Thr	Pro	Ser	Gln	Asn	Thr
				50					55					60	
Gln	Cys	Gln	Pro	Cys	Pro	Pro	Gly	Thr	Phe	Ser	Ala	Ser	Ser	Ser	Ser
				65					70					75	80
Ser	Glu	Gln	Cys	Gln	Pro	His	Arg	Asn	Cys	Thr	Ala	Leu	Gly	Leu	Ala
						85				90				95	
Leu	Asn	Val	Pro	Gly	Ser	Ser	Ser	His	Asp	Thr	Leu	Cys	Thr	Ser	Cys
				100						105				110	
Thr	Gly	Phe	Pro	Leu	Ser	Thr	Arg	Val	Pro	Gly	Ala	Glu	Glu	Cys	Glu
				115						120				125	
Arg	Ala	Val	Ile	Asp	Phe	Val	Ala	Phe	Gln	Asp	Ile	Ser	Ile	Lys	Arg
				130						135				140	
Leu	Gln	Arg	Leu	Leu	Gln	Ala	Leu	Glu							
				145						150					

SEQ ID NO:3:

```

      Met Ala Leu Lys Val Leu Pro Leu His Arg Thr Val Leu Phe Ala Ala
      1              5              10              15
Ile Leu Phe Leu Leu His Leu Ala Cys Lys Val Ser Cys Glu Thr Gly
      20              25              30
Asp Cys Arg Gln Gln Glu Phe Lys Asp Arg Ser Gly Asn Cys Val Leu
      35              40              45
Cys Lys Gln Cys Gly Pro Gly Met Glu Leu Ser Lys Glu Cys Gly Phe
      50              55              60
Gly Tyr Gly Glu Asp Ala Gln Cys Val Pro Cys Arg Pro His Arg Phe
      65              70              75              80
Lys Glu Asp Trp Gly Phe Gln Lys Cys Lys Pro Cys Ala Asp Cys Ala
      85              90              95
Leu Val Asn Arg Phe Gln Arg Ala Asn Cys Ser His Thr Ser Asp Ala
      100             105             110
Val Cys Gly Asp Cys Leu Pro Gly Phe Tyr Arg Lys Thr Lys Leu Val
      115             120             125
Gly Phe Gln Asp Met Glu Cys Val Pro Cys Gly Asp Pro Pro Pro Pro
      130             135             140
Tyr Glu Pro His Cys Thr Ser Lys Val Asn Leu Val Lys Ile Ser Ser
      145             150             155             160
      Thr Val Ser Ser Pro Arg Asp Thr Ala Val Ala Ala Val Ile Cys Ser
      165             170             175
Ala Leu Ala Thr Val Leu Leu Ala Cys Ser Ser Cys Val Ser Ser Thr
      180             185             190
Ala Arg Gly Ser Ser Trp Arg Arg Asn Pro Ala Val Ser Ser His Pro
      195             200             205
Ser Val
      210

```

SEQ ID NO:4:

```

      Met Ala Leu Lys Val Leu Pro Leu His Arg Thr Val Leu Phe Ala Ala
      1              5              10              15
Ile Leu Phe Leu Leu His Leu Ala Cys Lys Val Ser Cys Glu Thr Gly
      20              25              30
Asp Cys Arg Gln Gln Glu Phe Lys Asp Arg Ser Gly Asn Cys Val Leu

```

```

          35          40          45
Cys Lys Gln Cys Gly Pro Gly Met Glu Leu Ser Lys Glu Cys Gly Phe
          50          55          60
Gly Tyr Gly Glu Asp Ala Gln Cys Val Pro Cys Arg Pro His Arg Phe
          65          70          75          80
Lys Glu Asp Trp Gly Phe Gln Lys Cys Lys Pro Cys Ala Asp Cys Ala
          85          90          95
Leu Val Asn Arg Phe Gln Arg Ala Asn Cys Ser His Thr Ser Asp Ala
          100          105          110
Val Cys Gly Asp Cys Leu Pro Gly Phe Tyr Arg Lys Thr Lys Leu Val
          115          120          125
Gly Phe Gln Asp Met Glu Cys Val Pro Cys Gly Asp Pro Pro Pro Pro
          130          135          140
Tyr Glu Pro His Cys Glu
          145          150

```

SEQ ID NO 5:

```

          10          20          30          40          50
MVQLTQQTEL QSLRREVSRL QGTGGPSQNG EGYPWQSLPE QSSDALEAWE

```

```

          60          70          80          90
NGERSRKRRR VLTQKQKKQH SVLHLVPINA TSKDDSDVTE

```

```

          100          110          120          130          140          150
VMWQPALRRG RGLQAQGYGV RIQDAGVYLL YSQVLFQDVT FTMGQVVSRE QQGRQETLFR

```

```

          160          170          180          190          200
CIRSMPSHPD RAYNSCYSAG VFHLHQGDIL SVIIPRARAK LNLSPHGTFL GFVKL

```

SEQ ID NO: 6

ACCGGTCCGGAATTCCCGGGTCGACCCACGCGTCCGCCCACGCGTCCGAGAAGACTTTGAAATTCTTACAA
AAACTGAAAGTGAAATGAGGAAGACAGATTGAGCAATCCAATCGGAGGGTAAATGCCAGCAAACCTACTGT
ACAGTAGGGGTAGAGATGCAGAAAGGCAGAAAGGAGAAAATTCAGGATAACTCTCCTGAGGGGTGAGCCAA
GCCCTGCCATGTAGTGCACGCAGGACATCAACAAACACAGATAACAGGAAATGATCCATTCCCTGTGGTCA
CTTATTCTAAAGGCCCAACCTTCAAAGTTCAAGTAGTGATATGGATGACTCCACAGAAAGGGAGCAGTCA
CGCCTTACTTCTTGCCCTTAAGAAAAGAGAAGAAATGAAACTGAAGGAGTGTGTTTCCATCCTCCACGGAA
GGAAAGCCCCCTGTGTCGATCCTCCAAAGACGGAAAGCTGCTGGCTGCAACCTTGCTGCTGGCAGTGTGT
CTTGCTGCCTCACGGTGGTGTCTTTCTACCAGGTGGCCGCCCTGCAAGGGGACCTGGCCAGCCTCCGGGCA
GAGCTGCAGGGCCACCACGCGGAGAAGCTGCCAGCAGGAGCAGGAGCCCCCAAGCCGGCCTGGAGGAAGC
TCCAGCTGTCAACGCGGGACTGAAAATCTTTGAACCACAGCTCCAGGAGAAAGGCAACTCCAGTCAGAACA
GCAGAAATAAGCGTGCCGTTTCAGGGTCCAGAAGAAACAGTCACTCAAGACTGCTTGCAACTGATTGCAGAC
AGTGAAACACCAACTATACAAAAAGGATCTTACACATTTGTTCCATGGCTTCTCAGCTTTAAAGGGGAAG
TGCCCTAGAAGAAAAGAGAATAAAATATTGGTCAAAGAACTGGTTACTTTTTTATA
TATGGTCAGGTTTTATATACTGATAAGACCTACGCCATGGGACATCTAATTCAGAGGAAGAAGTCCATGT
CTTTGGGGATGAATTGAGTCTGGTGACTTTGTTTCGATGTATTCAAATATGCCTGAAACACTACCCAATA
ATTCTGCTATTTCAGCTGGCATTGCAAACTGGAAGAAGGAGATGAACTCCAATTGCAATACCAAGAGAA
AATGCACAAATATCACTGGATGGAGATGTCACATTTTTTGGTGCATTGAACTGCTGTGACCTACTTACAC
CATGTCTGTAGCTATTTTCTCCCTTTCTCTGTACCTCTAAGAAGAAAGAATCTAACTGAAAATAC
CAAAAAAAAAAAAAAAAAAAAAA

SEQ ID NO: 7:

CTGGAGCGCT GCCGCTACTG CAACGTCCTC TGCGGGGAGC GTGAGGAGGA GGCACGGGCT
60

TGCCACGCCA CCCACAACCG TGCCTGCCGC TGCCGCACCG GCTTCTTCGC GCACGCTGGT
120

TTCTGCTTGG AGCACGCATC GTGTCCACCT GGTGCCGGCG TGATTGCCCC GGGCACCCCC
180

AGCCAGAACA CGCAGTGCCA GCCGTGCCCC CCAGGCACCT TCTCAGCCAG CAGCTCCAGC
240

TCAGAGCAGT GCCAGCCCCA CCGCAACTGC ACGGCCCTGG GCCTGGCCCT CAATGTGCCA
300

GGCTCTTCCT CCCATGACAC CCTGTGCACC AGCTGCACTG GCTTCCCCCT CAGCACCAGG
360

GTACCAGGAG CTGAGGAGTG TGAGCGTGCC GTCATCGACT TTGTGGCTTT CCAGGACATC
420

TCCATCAAGA GGCTGCAGCG GCTGCTGCAG GCCCTCGAG
459

SEQ ID NO:8:

TCCGGCGCCG CGGGGCAGGA CAAGGGAAG GAATAACAC GTTTGGTGAG AGCCATGGCA
60

CTCAAGGTCC TACCTCTACA CAGGACGGTG CTCTTCGCTG CCATTCTCTT CCTACTCCAC
120

CTGGCATGTA AAGTGAGTTG CGAAACCGGA GATTGCAGGC AGCAGGAATT CAAGGATCGA
180

TCTGGAAACT GTGTCCTCTG CAAACAGTGC GGACCTGGCA TGGAGTTGTC CAAGGAATGT
240

GGCTTCGGCT ATGGGGAGGA TGCACAGTGT GTGCCCTGCA GGCCGCACCG GTTCAAGGAA
300

GACTGGGGTT TCCAGAAGTG TAAGCCATGT GCGGACTGTG CGCTGGTGAA CCGCTTTCAG
360

AGGGCCAACT GCTCACACAC CAGTGATGCT GTCTGCGGGG ACTGCCTGCC AGGATTTTAC
420

CGGAAGACCA AACTGGTTGG TTTTCAAGAC ATGGAGTGTG TGCCCTGCGG AGACCCACCT
480

CCTCCCTACG AACCACACTG TACCAGCAAG GTGAACCTTG TGAAGATCTC CTCCACCGTC
540

TCCAGCCCTC GGGACACGGC GGTGGCTGCC GTCATCTGCA GTGCTCTGGC CACGGTGCTG
600

CTCGCCTGCT CATCCTGTGT GTCATCTACT GCAAGAGGCA GTTCATGGAG AAGAAACCCA
660

GCTGTAAGCT CCCATCCCTC TGTCTCACTG TGAAGTGAGC TTGTTAGCAT TGTCAACCAA
720

GAGTTCTCAA GACACCTGGC TGAGACCTAA GACCTTTAGA GCATCAACAG CTACTTAGAA
780

TACAAGATGC AGGAAAACGA GCCTCTTCAG GAATCTCAGG GCCTCCTAGG GATGCTGGCA
840

AGGCTGTGAT GTCTCAAGGC TACCAGGAAA AAATAAAAGT TGTCTATACC CTA
893

SEQ ID NO:9:

GAGGCAAGAT TCGGCACGAG GCGGTTTGGC GCGGAAGTGC TACCAAGCTG CGGAAAGCGT
60

GAGTCTGGAG CACAGCACTG GCGAGTAGCA GGAATAAACA CGTTTGGTGA GAGCCATGGC
120

ACTCAAGGTC CTACCTCTAC ACAGGACGGT GCTCTTCGCT GCCATTCTCT TCCTACTCCA
180

CCTGGCATGT AAAGTGAGTT GCGAAACCGG AGATTGCAGG CAGCAGGAAT TCAAGGATCG
240

ATCTGGAAAC TGTGTCCTCT GCAAACAGTG CGGACCTGGC ATGGAGTTGT CCAAGGAATG
300

TGGCTTCGGC TATGGGGAGG ATGCACAGTG TGTGCCCTGC AGGCCGCACC GGTTC AAGGA
360

AGACTGGGGT TTCCAGAAGT GTAAGCCATG TGCGGACTGT GCGCTGGTGA ACCGCTTTCA
420

GAGGGCCAAC TGCTCACACA CCAGTGATGC TGTCTGCGGG GACTGCCTGC CAGGATTTTA
480

CCGGAAGACC AACTGGTTG GTTTTCAAGA CATGGAGTGT GTGCCCTGCG GAGACCCACC
540

TCCTCCCTAC GAACCACACT GTGAGTGATG TGCCAAGTGG CAGCAGACCT TTAAAAAAAA
600

AAGAAAAAAAA AACAAACAAA AAC
623

SEQ ID NO 10:

10 20 30 40 50 60 70 80 90
CTTCCTAGAG GGACTGGAAC CTAATTCTCC TGAGGCTGAG GGAGGGTGGG GGGTCTCAAG GCAACGCTGG CCCCACGACG GAGTGCCAGG
GAAGGATCTC CCTGACCTTG GATTAAGAGG ACTCCGACTC CCTCCCACTC CCCAGAGTTC CGTTGCGACC GGGGTGCTGC CTCACGGTCC

100 110 120 130 140 150 160 170 180
AGCACTAACA GTACCCCTAG CTTGCTTTCC TCCTCCCTCC TTTTATTTT CAAGTTCCTT TTTATTTCTC CTTGCGTAAC AACCTTCTTC
TCGTGATTGT CATGGGAATC GAACGAAAGG AGGAGGGAGG AAAAAATAAA GTTCAAGGAA AAATAAAGAG GAACGCATTG TTGGAAGAAG

190 200 210 220 230 240 250 260 270
CCTTCTGCAC CACTGCCCGT ACCCTTACCC GCCCCGCCAC CTCCTTGCTA CCCCCTCTT GAAACCACAG CTGTTGGCAG GGTCCCCAGC
GGAAGACGTG GTGACGGGCA TGGGAATGGG CGGGGCGGTG GAGGAACGAT GGGGTGAGAA CTTTGGTGTC GACAACCGTC CCAGGGGTGC

280 290 300 310 320 330 340 350 360
TCATGCCAGC CTCATCTCCT TTCTTGCTAG CCCCCAAGS SCTCCAGGCA ACATGGGGGG CCCAGTCAAG GAGCCCGCAC TCTCAGTTGC
AGTAGCGTCG GAGTAGAGGA AAGAAGCATC GGGGTTCCTC SGAGGTCCGT TGTACCCCCC GGGTCAGTCT CTCGGCCGTG AGAGTCAACG

370 380 390 400 410 420 430 440 450
CCTCTGGTTG AGTTGGGGGG CAGCTCTGGG GCGCGTGSBT TGTGCATGGT YBAGCTGACC CAACAAACAG AGCTGCAGAS CCTCAGGAGA
GGAGACCAAC TCAACCCCCC GTCGAGACCC CCGGCACSV A ACACGTACCA RVTGCACTGG GTTGTGTTGC TCGACGTCTC GGAGTCTCTC

460 470 480 490 500 510 520 530 540
GAGGTGAGCC GGCTGCAGGG GACAGGAGGC CCCCCTCCAGA ATGGGGAAGG GTATCCCTGG CAGAGTCTCC CGGAGCAGAG TTCCGATGCC
CTCCACTCGG CCGACGTCCT CTGTCTCTCG GGGAGGGTCT TACCCCTTCC CATAGGGACC GTCTCAGAGG GCCTCGTCTC AAGGCTACGG

550 560 570 580 590 600 610 620 630
CTGGAAGCCT GGGAGAATGG GGAGAGATCC CGGAAAAGGA GAGCAGTGCT CACCCAAAAA CAGAAGAAGC AGCACTCTGT CCTGCACCTG
GACCTTCGGA CCTCTTACC CCTCTTAGG GCCTTTTCCT CTCGTACAGA GTGGGTTTTGT GTCTTCTTCC TCGTGAGACA GGACGTGGAC

640 650 660 670 680 690 700 710 720
GTTCCCATTA ACGCCACCTC CAAGGATGAC TCCGATGTGA CAGAGGTGAT GTGGCAACCA GCTCTTAGGC GTGGGAGAGG CCTACAGGCC
CAAGGGTAAT TGCGGTGGAG GTTCTACTG AGGCTACACT GTCTCCACTA CACCGTTGGT CGAGAATCCG CACCCCTCTC GGATGTCCGG

730 740 750 760 770 780 790 800 810
CAAGGATATG GTGTCCGAAT CCAGGATGCT GGAGTTTATC TGCTGTATAG CCAGGTCTCT TTTCAAGACG TGACTTTTAC CATGGGTCCG
GTTCTTATAC CACAGGCTTA GGTCTTACGA CCTCAAATAG ACGACATATC GGTCCAGGAC AAAGTTCTCG ACTGAAAGTG GTACCCAGTC

820 830 840 850 860 870 880 890 900
GTGGTGTCTC GAGAAAGGCA AGGAAAGGCA GAGACTCTAT TCCGATGTAT AAGAAGTATG CCTTCCAC CCGACCGGGC CTACAACAGC
CACCACAGAG CTCTTCCGGT TCCTTCCGTC CTCTGAGATA AGGCTACATA TTCTTCATAC GGGAGGGTGG GCCTGGCCCG GATGTTGTCC

910 920 930 940 950 960 970 980 990
TGCTATAGCG CAGGTGTCTT CCATTTACAC CAAGGGGATA TTCTGAGTGT CATAATTCCC CGGGCAAGGG CGAAACTTAA CCTCTCTCCA
ACGATATCCG GTCCACAGAA GGTAAATGTG GTTCCCTTAT AAGACTCACA GTATTAAAGG GCCCGTTCCT GCTTTGAATT GGAGAGAGGT

1000 1010 1020 1030 1040 1050 1060 1070 1080
CATGGAACCT TCCTGGGGTT TGTGAAACTG TGATTGTGTT ATAAAAAGTG GCTCCAGCT TGGAAGACCA GGGTGGGTAC ATACTGGAGA
GTACCTTGA AGGACCCCAA ACACCTTGAC ACTAACACAA TATTTTTCAC CGAGGGTCCA ACCTTCTGTT CCCACCATG TATGACCTCT

1090 1100 1110 1120 1130 1140 1150 1160 1170
CAGCCAAGAG CTGAGTATAT AAAGGAGAGG GAATGTGCAG GAACAGAGGC ATCTTCTCTG GTTTGGCTCC CCGTTCTCTA CTTTCTCCCT
GTCGGTCTC GACTCATATA TTTCTCTCTC CTTACACGTC CTTGTCTCCG TAGAAGGACC CAAACCGAGG GGCAAGGAGT GAAAAGGGAA

1180 1190 1200 1210 1220 1230 1240 1250 1260
TTCAATCCCA CCCCCTAGAC TTTGATTTTA CGGATATCTT GCTTCTGTTT CCGATGGAGC TCCGAATTCT TGCGTGTGTG TAGATGAGGG
AAGTAAGGGT GGGGATCTG AACTAAAAAT GCCTATAGAA CGAAGACAAG GGTACCTCG AGGCTTAAGA ACGCACACAC ATCTACTCCC

SEQ ID NO:11
AGGTCCATGTCTTTGGG

SEQ ID NO:12
GGGGATGAATTGAGTCTG

SEQ ID NO:13
GTGCCCTCGAAGAAAAAG

SEQ ID NO:14
GCAAGTTGGAGTTCATC

SEQ ID NO:15
EYMPTD

SEQ ID NO:16
EYMPTE



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, 15/19, C07K 14/715, 14/52, C12Q 1/68, G01N 33/53, 33/68, C07K 16/24, 16/28	A3	(11) International Publication Number: WO 99/33980 (43) International Publication Date: 8 July 1999 (08.07.99)
(21) International Application Number: PCT/US98/27474 (22) International Filing Date: 22 December 1998 (22.12.98) (30) Priority Data: 60/068,959 30 December 1997 (30.12.97) US 09/212,270 16 December 1998 (16.12.98) US (71) Applicant: CHIRON CORPORATION [US/US]; 4560 Horton Street -R440, Emeryville, CA 94608 (US). (72) Inventors: TRIBOULEY, Catherine; Chiron Corporation, In- tellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US). POT, David; Chiron Corporation, Intel- lectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US). KASSAM, Altaf; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US). LAMSON, George; Chiron Cor- poration, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US). (74) Agents: POTTER, Jane, E., R. et al.; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims</i> <i>and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 18 November 1999 (18.11.99)
(54) Title: MEMBERS OF TNF AND TNFR FAMILIES		
(57) Abstract Members of the TNF and the TNFR superfamily of proteins have been identified. These proteins are promising targets for therapeutic intervention and mimesis. TNF-L and TNFR-L proteins can be used to induce cell death and/or proliferation of cells. Members of these superfamilies have been implicated in a broad variety of disease processes, making them central biological and physiological regulators.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/27474

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C12N15/19 C07K14/715 C07K14/52 C12Q1/68
G01N33/53 G01N33/68 C07K16/24 C07K16/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K C12Q G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	WO 99 12964 A (BIOGEN INC. (US); TSCHOPP JURG) 18 March 1999 (1999-03-18) abstract page 14, line 7-20 page 23, line 10-19 page 28 -page 31; claims Seq.ID:1,2 page 32 figure 1 --- -/--	1-16

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *B* document member of the same patent family

Date of the actual completion of the international search

27 April 1999

Date of mailing of the international search report

24.09.99

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Macchia, G

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/27474

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 98 18921 A (HUMAN GENOME SCIENCES INC.; YU GUO-LIANG; EBNER REINHARD; NI JIAN) 7 May 1998 (1998-05-07) abstract page 11, line 21 -page 12, line 11 page 41, line 19 -page 42, line 2 page 54 -page 57 Seq.ID:1,2 page 77 -page 80 page 90 -page 94; claims figures 1,2 ---	1-16
P,X	WO 98 27114 A (SCHERING CORPORATION (US); GORMAN DANIEL M.) 25 June 1998 (1998-06-25) abstract page 39, line 18-32 page 47, line 6-20 Seq.ID:3,4 page 58 -page 61 page 67 -page 68; claims ---	1-16
P,X	EP 0 869 180 A (SMITHKLINE BEECHAM CORPORATION (US); HURLE MARK R.; YOUNG PETER R.) 7 October 1998 (1998-10-07) the whole document ---	1-16
E	WO 99 11791 A (UNIVERSITY OF WASHINGTON (US); CHAUDHARY PREET M.) 11 March 1999 (1999-03-11) abstract page 30, line 3 -page 34, line 32 page 54, line 10-25 page 57, line 1 -page 59, line 12 page 129 -page 130; claims 33-38 figure 11A ---	1-16
P,X	WO 98 55620 A (REGENERON PHARMACEUTICALS (US); MASIAKOWSKI PIOTR; VALENZUELA DAVID) 10 December 1998 (1998-12-10) the whole document ---	1-16
P,X	WO 98 55621 A (REGENERON PHARMACEUTICALS (US); MASIAKOWSKI PIOTR; VALENZUELA DAVID) 10 December 1998 (1998-12-10) the whole document ---	1-16
X	Database EMBL, ID AA682496 Accession number AA682496 10 December 1997 99% identity with Seq.ID:6 nt.807-1260. reverse orientation XP002101204 the whole document ---	9

	-/--	

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/27474

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>GRUSS H.-J. AND DOWER S.K.: "Tumor necrosis factor ligand superfamily: involvement in the pathology of malignant lymphomas"</p> <p>BLOOD, vol. 85, no. 12, 15 June 1995 (1995-06-15), pages 3378-3404, XP002094502</p> <p style="text-align: center;">---</p>	
A	<p>AGGARWAL B.B. AND NATARAJAN K.: "Tumor necrosis factor: developments during the last decade"</p> <p>EUROPEAN CYTOKINE NETWORK, vol. 7, no. 2, April 1996 (1996-04) - June 1996 (1996-06), pages 93-124, XP002094503</p> <p style="text-align: center;">---</p>	
A	<p>GRUSS H.-J.: "Molecular, structural, and biological characteristics of the tumor necrosis factor ligand superfamily"</p> <p>INTERNATIONAL JOURNAL OF CLINICAL AND LABORATORY RESEARCH, vol. 26, no. 3, 1996, pages 143-159, XP002094504</p> <p style="text-align: center;">---</p>	
A	<p>WO 97 33902 A (HUMAN GENOME SCIENCES INC. (US); NI JAN; YU GUO-LIANG; GENTZ REINER L.) 18 September 1997 (1997-09-18)</p> <p style="text-align: center;">-----</p>	

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/ 27474

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see ADDITIONAL SHEET

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-16 all partially

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-16 all partially

Protein having an amino acid sequence as in Seq.ID:1, variants, fusion proteins thereof. Antibody binding to said protein. cDNA molecule encoding said protein, as in Seq.ID:6, homologous sequences thereof. Construct comprising said polynucleotide, host cell comprising said construct. Method of screening for a compound capable of modulating cell death, involving said protein. Method of identifying a binding partner of said protein.

2. Claims: 1-16 all partially

As invention 1 but concerning Seq.ID:2 and 7.

3. Claims: 13-16 all partially

Method of screening for a compound capable of modulating cell death, involving a protein comprising an amino acid sequence as in Seq.ID:3,4 encoded by a nucleotide sequence as in Seq.ID:8,9. Method of identifying a binding partner of said protein.

4. Claims: 13-16 all partially

As invention 3 but concerning Seq.ID:5 and 10.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/27474

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9912964 A	18-03-1999	AU 9315298 A AU 9316298 A WO 9912965 A	29-03-1999 29-03-1999 18-03-1999
WO 9818921 A	07-05-1998	AU 7674596 A EP 0939804 A	22-05-1998 08-09-1999
WO 9827114 A	25-06-1998	AU 5705898 A	15-07-1998
EP 0869180 A	07-10-1998	CA 2232743 A JP 10323194 A	02-10-1998 08-12-1998
WO 9911791 A	11-03-1999	AU 9376498 A	22-03-1999
WO 9855620 A	10-12-1998	AU 7608898 A AU 7713098 A WO 9855621 A	21-12-1998 21-12-1998 10-12-1998
WO 9855621 A	10-12-1998	AU 7608898 A AU 7713098 A WO 9855620 A	21-12-1998 21-12-1998 10-12-1998
WO 9733902 A	18-09-1997	AU 5366596 A EP 0897390 A	01-10-1997 24-02-1999